



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : C12N 15/82, 15/11	A2	(11) International Publication Number: WO 00/63397 (43) International Publication Date: 26 October 2000 (26.10.00)
(21) International Application Number: PCT/EP00/03521 (22) International Filing Date: 17 April 2000 (17.04.00) (30) Priority Data: 09/294,022 20 April 1999 (20.04.99) US (71) Applicant: AVENTIS CROPS SCIENCE N.V. [BE/BE]; Jozef Plateastraat 22, B-9000 Gent (BE). (72) Inventors: MEULEWAETER, Frank; Wechaagstraat 73, B-9160 Eksaarde (BE). CORNELISSEN, Marc; Ellebogten 38, B-9070 Heusden (BE). JACOBS, John; Wilgenstraat 4, B-9820 Merelbeke (BE). VAN ELDIK, Gerben; Nekkersberglaan 41, B-9000 Gent (BE). METZLAF, Michael; Irislaan 26, B-3080 Tervuren (BE).		(81) Designated States: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHODS AND MEANS FOR DELIVERING INHIBITORY RNA TO PLANTS AND APPLICATIONS THEREOF <div style="text-align: center;"> <p><u>Tobacco mosaic virus</u></p> </div> (57) Abstract <p>The invention provides methods and means for the identification of genes involved in the determination of a plant trait or for the identification encoded by a nucleic acid comprising a determined nucleotide sequence. The invention also provides kits comprising viral RNA vectors derived from satellite viruses and corresponding helper viruses for the introduction of inhibitory RNA into plant cells and plants.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Methods and means for delivering inhibitory RNA to plants and applications thereof

Field of the invention:

5 This invention relates to the field of functional genomics in plants, more particularly it relates to methods for the further identification and isolation of a nucleic acid with a nucleotide sequence of interest in a collection of preselected nucleic acid sequences correlated with a particular trait, preferably an agronomical important trait, using a kit of viral RNA vectors
10 which allow systemic spread of all components of the kit in a plant, wherein one of the viral RNA vectors comprises a library of gene-silencing constructs for the preselected nucleic acid sequences. The invention also relates to a method for modulating, preferably reducing, particularly eliminating the expression of a selected nucleic acid sequence, using the viral vector kit,
15 whereby one of the vectors comprises a gene-silencing construct for the selected nucleic acid sequence. The latter method may be used for validating the function of a nucleic acid sequence whose expression is correlated with the presence or absence of a specific trait in plants, but with otherwise unknown function. Preferably, one of the viral RNA vector components of the
20 kit is a vector derived from a satellite virus.

Background art

25 The recent, rapid expansion of available nucleic acid sequence information has necessitated the development of methods for identifying the function of nucleic acid sequences, particularly transcribed nucleic acid sequences such as expressed sequence tags, with unknown function, in an efficient and labor-cost effective way.

30

To identify the role of sequenced nucleic acids from plants of unknown function it is necessary to produce or identify plants in which those nucleic acids are either structurally or functionally inactivated. Plants wherein

CONFIRMATION COPY

expression of both sense and antisense constructs in cells of one plant. The sense and antisense nucleic acids may be in the same transcriptional unit, so that a single RNA transcript that has self-complementarity is generated upon transcription.

5

Hamilton et al. (1998) describe improved silencing e.g. of tomato ACC-oxidase gene expression using a sense RNA containing two additional upstream inverted copies of its 5' untranslated region.

10

WO 98/53083 describes constructs and methods for enhancing the inhibition of a target gene within an organism, involving the insertion into the gene silencing vector of an inverted repeat of all or part of a polynucleotide region within the vector.

15

It should be clear however, that the use of inhibitory RNA as a tool in reversed genetics analysis of gene function via high throughput methods, whereby the inhibitory RNA is generated from gene-silencing constructs which are stably integrated in the genome of transgenic plants, suffers from the same drawbacks as the methods wherein the nucleotide sequences are structurally

20

inactivated.

EP 0 194 809 and US 5,500,360 suggest the use of viral RNA vectors to produce regulatory RNA such as anti-sense RNA.

25

Initial exploration of the use of viral vectors to deliver inhibitory RNA into cells of plants has been described by Chapman (1991). In this publication, gene silencing constructs comprising nucleotide sequences complementary to the translated region of the GUS gene on a PVX derived viral vector were described. The experiments however, remained inconclusive as to whether

30

gene silencing could be provoked using viral vectors for the production of inhibitory RNA.

WO 93/03161 is directed toward recombinant plant viral nucleic acids and to hosts infected thereby. The non-native nucleic acid sequence which is transcribed may be transcribed as an RNA which is capable of regulating the expression of a phenotypic trait by an anti-sense mechanism.

5

English et al., 1996 describe the suppression of the accumulation of a viral vector comprising a foreign nucleotide sequence in transgenic plants exhibiting silencing of nuclear genes comprising the same foreign nucleotide sequences, thus linking gene silencing and viral vectors, albeit in a reverse way as envisioned here.

10

Kumagai et al. 1995 (PNAS 92, 1679-1683) described the inhibition of phytoene desaturase gene by viral delivery of antisense RNA.

15

WO 95/34668 suggests the use of genetic constructs based on RNA viruses which replicate in the cytoplasm of cells to provide inhibitory RNA, either antisense or co-suppressor (sense) RNA.

20

Baulcombe et al. (1998) and Ruiz et al. (1998) describe virus-induced gene silencing of the endogenous phytoene desaturase gene (PDS) or of a green fluorescent protein transgene (GFP) in plants, using potato virus X derived vectors carrying inserts homologous to PDS and GFP, respectively. The authors further suggested that virus-induced gene silencing may develop into a novel assay of gene function, by introducing a fragment of the genome of a viral vector and inferring the function of the gene from the symptoms of the infected plants exhibiting gene silencing.

25

The described methods for identification of the function of a gene with known nucleotide sequence however, have drawbacks and limitations. In the first place, the applicability of the mentioned viral RNA vector based gene silencing methods on larger scale is in practice limited to the identification of genes with essential functions or genes with macroscopically visible phenotypes. Secondly, all methods employ viral vectors which are capable of

30

autonomous replication in plant cells and cell-to-cell movement , whereby care has to be taken not to inactivate the essential functions required for these functions. This may particularly be a disadvantage when tailoring these methods to the needs of particular plants, such as crop plants, by developing
5 new viral vectors more apt for replication and systemic spread in the plants.

The prior art is thus deficient in the lack of efficient methods for large scale identification of the function of nucleic acids with known nucleotide sequence, or for the isolation of the genes of interest from a pool of genes with known
10 nucleotide sequence, but unknown function.

Brief Description of the figures

15 Figure 1 is a schematic representation of the viral RNA vectors used in the Examples. MP: movement protein; CP: coat protein; OAS: origin of assembly. The open reading frames are indicated by boxes. Each original viral genome is characterized by a specific pattern.

20 ■■■ : tobacco mosaic virus; ■■■■ tobacco necrosis virus; ■■■■ satellite tobacco mosaic virus; ■■■■ satellite tobacco necrosis virus.

Summary of the invention

The invention provides a method for isolating genes involved in the
25 determination of a trait or a phenotype of a plant species, comprising identifying a set of nucleic acids sequences of genes, whose expression is correlated with a trait of interest; creating a library of gene silencing constructs targeted or adapted to the nucleotide sequence of the identified nucleic acids in a viral RNA vector which is capable of replication inside plant cells and
30 optionally, movement between plant cells of a plant; infecting a collection of individual plants of the same plant species with the library of gene silencing constructs whereby each plant is infected with at least one member of the library; identifying a plant wherein the trait or phenotype is altered using an assay adapted to that trait or phenotype; and isolating the gene involved in the
35 determination of the trait or phenotype in the plant species, from the library, based on the nucleotide sequence to which the gene silencing construct in the

identified plant was targeted. Preferably the viral RNA vector is capable of autonomous replication inside plant cells and optionally autonomous movement between plant cells and particularly the viral RNA vector is derived from cowpea mosaic virus.

5

Alternatively, the viral RNA vector, is derived from a satellite RNA virus, preferably satellite tobacco mosaic virus or satellite tobacco necrosis virus, particularly it further comprises an origin of assembly of tobacco mosaic virus, and is capable of replication inside plant cells and optionally movement
10 between plant cells when the required factors are supplemented *in trans*, preferably by infection with a helper virus, preferably tobacco mosaic virus or a helper virus which is derived from tobacco necrosis virus and comprises a gene encoding a coat protein gene of tobacco mosaic virus, and optionally, a movement protein of tobacco mosaic virus.

15

Gene-silencing constructs comprised within the viral RNA vector may comprise antisense RNA or sense RNA, preferably they may comprise an inverted repeat. Particularly the gene-silencing constructs comprise complementary stretch of at least 50, preferably at least 100 nucleotides of
20 sense and antisense RNA. Especially preferred are gene-silencing constructs comprising at least two copies of part of the nucleotide sequences of the collection of nucleic acids, the copies being in inverted repeat.

25

It is another object of the invention to provide a method for the isolation or selection of a nucleic acid with a specific function from a collection of nucleic acids, wherein the collection of nucleic acids is characterized by the fact that variation in the expression pattern of the nucleic acids is correlated with variation in a trait or phenotype of a plant harboring the nucleic acids comprising the steps of creating a library of gene silencing constructs targeted
30 or adapted to the nucleotide sequence of the identified nucleic acids in a viral RNA vector which is capable of replication inside and optionally movement between plant cells; infecting a collection of plants with the library of gene silencing constructs, whereby each plant is infected with one member of the

library; identifying plants with altered trait or phenotype using an assay adapted to the trait or phenotype under investigation; and optionally isolating the nucleic acid with the specific function from the identified plant with altered trait or phenotype.

5

It is yet another object of the invention to provide a method for determining the function encoded by a nucleic acid comprising a known nucleotide sequence in a plant, comprising the steps of providing a viral RNA vector derived from a satellite RNA virus, comprising a gene-silencing construct targeted to a gene
10 comprising the known nucleotide sequence; infecting or inoculating the plant with the chimaeric viral RNA vector and a corresponding helper virus or helper virus RNA; and identifying an altered trait or phenotype of the co-infected plant.

15

The invention further provides a method for isolating essential genes from a plant, comprising creating a library of random gene-silencing constructs, preferably by cloning random DNA fragments of the plant or by cloning random cDNA fragments, particularly duplicated cDNA fragments in inverted repeat in a cDNA copy of the viral RNA vector derived from a satellite RNA
20 virus, preferably STMV or STNV, particularly a viral RNA vector comprising an origin of assembly of TMV; infecting a plant with individual members of the library and with a corresponding helper virus, preferably tobacco mosaic virus or a helper virus derived from tobacco necrosis virus and comprising the coat protein of tobacco mosaic virus and optionally the movement protein;
25 identifying plants developing gene-silencing-construct-associated necrosis and optionally isolating the viral RNA vector from the necrotized tissue.

30

It is yet another object of the invention to provide a method for introduction of inhibitory RNA, preferably sense or antisense RNA, particularly inhibitory RNA comprising an inverted repeat, especially inhibitory RNA comprising a
complementary stretch of at least 50, preferably at least 100 nucleotides of sense and antisense RNA, into plant cells, preferably into the cytoplasm of plant cells, comprising introducing into a plant cell, a viral RNA vector

comprising the inhibitory RNA or comprising a chimeric nucleic acid which when transcribed yields the inhibitory RNA, wherein the viral RNA vector is derived from a satellite RNA virus, preferably STMV or STNV, particularly a STMV - derived or STNV derived RNA vector comprising an origin of assembly from tobacco mosaic virus; and introducing into the same plant cell, a corresponding helper virus, preferably tobacco mosaic virus or a chimaeric helper virus derived from tobacco necrosis virus and comprising a coat protein gene of tobacco mosaic virus and optionally, a movement protein encoding gene of tobacco mosaic virus.

10

The invention further provides a kit for introduction of inhibitory RNA, preferably sense or antisense RNA, particularly inhibitory RNA comprising an inverted repeat, especially inhibitory RNA comprising a complementary stretch of at least 50, preferably at least 100 nucleotides of sense and antisense RNA in the cytoplasm of a plant cell comprising 1) a viral RNA vector derived from a satellite RNA virus, comprising the inhibitory RNA or which comprise a chimeric nucleic acid which when transcribed yields the inhibitory RNA; and 2) a corresponding helper virus.

20

A particularly preferred kit comprises 1) a viral RNA vector derived from a satellite tobacco mosaic virus, comprising an origin of assembly of tobacco mosaic virus, further comprising or encoding the inhibitory RNA; and 2) a corresponding helper virus derived from a tobacco mosaic virus.

25

Another particularly preferred kit comprises 1) a viral RNA vector derived from satellite tobacco necrosis virus, especially STNV-2 or STNV-C and comprising an origin of assembly of tobacco mosaic virus further comprising or encoding the inhibitory RNA; and 2) a corresponding helper virus derived from tobacco necrosis virus, particularly TNV-A or TNV-D, which comprises a coat protein gene of tobacco mosaic virus and optionally a movement protein gene of tobacco mosaic virus.

30

Detailed description of preferred embodiments.

The following definitions apply throughout this application, unless otherwise specified.

5 As used herein "comprising" is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, e.g., a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more
10 nucleotides or amino acids than the actually cited ones, i.e., be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA region which is functionally or structurally defined, may comprise additional DNA regions etc.

15 As used herein, "a trait of a plant" indicates a phenotype which is the combined result of the coordinated expression of a number of genes. Typical traits include yield, heterosis, drought-resistance, stress-resistance, high or low temperature-resistance, vigor, seed yield, plant habitat, architecture etc. Typically, a trait of a plant is named after its intended appearance.

20 A used herein a "phenotype" of a plant refers to any quantitative or qualitative characteristic of that plant, be it morphological (including macroscopic and microscopic characteristics), biochemical (including the presence, absence or concentration of particular metabolites or molecules) functional or other.

25 The term "gene" means any DNA or RNA fragment comprising a region (the "transcribed region") which is transcribed into a RNA molecule (e.g., a mRNA) in a cell, operably linked to suitable regulatory regions, e.g., a plant-expressible promoter. A gene may thus comprise several operably linked
30 fragments such as a promoter, a 5' leader sequence, a coding region, and a 3' region comprising a polyadenylation site. A plant gene endogenous to a particular plant species or virus (endogenous plant or virus gene) is a gene which is naturally found in that plant species or virus, or which can be introduced in that plant species by breeding techniques such as conventional
35 breeding techniques. A chimeric gene is any gene which is not normally found

in a plant species or, alternatively, any gene in which the promoter is not associated in nature with part or all of the transcribed DNA region or with at least one other regulatory region of the gene.

5 The term "expression of a gene" refers to the process wherein a DNA or RNA region which is operably linked to appropriate regulatory regions, particularly to a promoter, is transcribed into an RNA which is biologically active i.e., which is either capable of interaction with another nucleic acid or which is capable of being translated into a biologically active polypeptide or protein. A
10 gene is said to encode an RNA when the end product of the expression of the gene is biologically active RNA, such as e.g. an antisense RNA, a ribozyme or a replicative intermediate. A gene is said to encode a protein when the end product of the expression of the gene is a biologically active protein or polypeptide. In addition to the above defined elements, a gene may further
15 comprise elements for cap-independent translation such as an internal ribosome entry sequence or the first and second translation enhancing elements as defined in WO 97/49814.

As used herein the terms "gene-silencing" or "inhibitory" are not to be
20 interpreted as meaning a complete abolishing of the expression of the target gene(s) but also includes any reduction in expression, measured either as a reduction in transcription and/or translation, as a reduction in the accumulation of transcripts or translation products such as proteins, or as a reduction in the phenotypic expression of the target gene.

25 The term "reduction of phenotypic expression" refers to the comparison of the phenotypic expression of the nucleic acid of interest in the eukaryotic cell in the presence of the inhibitory RNA or gene-silencing constructs of the invention, to the phenotypic expression of the nucleic acid of interest in a
30 similar eukaryotic cell in the absence of the inhibitory RNA or gene-silencing constructs of the invention. The phenotypic expression in the presence of the inhibitory RNA of the invention should thus be lower than the phenotypic expression in absence thereof, preferably be only about 25%, particularly only

about 10%, more particularly only about 5% of the phenotypic expression in absence of the inhibitory RNA, especially the phenotypic expression should be completely inhibited for all practical purposes by the presence of the inhibitory RNA or the gene-silencing construct encoding such an RNA.

5

A reduction of phenotypic expression of a nucleic acid where the phenotype is a qualitative trait means that in the presence of the inhibitory RNA, the phenotypic trait switches to a different discrete state when compared to a situation in which such inhibitory RNA is absent. A reduction of phenotypic
10 expression of a nucleic acid may thus a.o. be measured as a reduction in transcription of (part of) that nucleic acid or reduction in the level of transcript, a reduction in translation of (part of) that nucleic acid or reduction in the level of translation products, or a reduction of the effect the presence of the transcribed RNA(s) or translated polypeptide(s) have on the eucaryotic cell or
15 the organism, and will ultimately lead to altered phenotypes. It is clear that the reduction in phenotypic expression of a nucleic acid of interest, may be accompanied by or correlated to an increase in a phenotype or trait.

In one embodiment of the invention, a method is provided to identify and
20 isolate genes involved in the determination of a trait or a phenotype of a plant. To this end, nucleic acid sequences are identified whose expression is correlated with the trait and/or phenotype of interest. Methods and means are available in the art for the almost simultaneous identification and/or isolation of a large number, if not the predominant part, of nucleotide sequences whose
25 expression, particularly whose transcription, is influenced subsequent to a stimulus corresponding to the trait to be investigated, in comparison with expression of these nucleotide sequences in a control plant. Such methods include but are not limited to differential display methods, such as the gel - based RNA differential display methods described by (Prahasar et al. 1996)
30 As a result of these methods, a collection of at least partially characterized nucleotide sequences with altered expression in response to a particular stimulus is identified. Typically, however, the application of such methods does not allow to discriminate between genes whose altered expression is

directly caused by the stimulus, and those who are further downstream in the chain of events and are only indirectly influenced by the stimulus and a further selection amongst the obtained collection of nucleotide sequences will be required. Even less do these methods allow to predict whether the inverse relationship also holds, i.e. whether influencing the expression of genes with particular nucleotide sequences, identified in the above mentioned way, also influences the trait of interest. A further validation of the obtained sequences is thus required, and preferably one which immediately verifies the above mentioned inverse relationship. To achieve this goal in a efficient and cost-effective way, a library of gene-silencing constructs may be created in a viral RNA vector which is capable of replication inside plant cells and movement between cells of a plant, so as to assure an efficient systemic spread within a plant which has been infected with a clone of such a library. The created library of gene-silencing constructs comprised within a viral RNA vector is then used to infect a representative number of plants in such a way that at each plant is infected by at least one member (one clone) of the library. The infected plants can then be analyzed to identify those plants which exhibit alterations in the trait under investigation, using an assay which is adapted to the trait under investigation. It is clear that this fine-tuning of assay and trait under investigation may be an important advantage over the existing methods for high throughput analysis, particularly when analyzing traits and/or phenotypes which do not result in macroscopically visible alterations, such as but not limited to modifications in specific metabolic pathways or alterations which are only detectable under specific conditions (e.g. heat, stress, drought-tolerance, pathogen-infection, application of specific herbicides or insecticides etc.).

The subset of nucleic acid sequences may also be identified on the basis of the presence of a particular signature characteristic of a class of proteins, such as but not limited to a kinase-specific domain, a binding motif etc.

The gene-silencing construct may then be isolated from the library or from the plant exhibiting the altered trait or phenotype, and be used to isolate the

corresponding gene based on the nucleotide sequence towards which the gene silencing construct was targeted.

In a preferred embodiment, the viral RNA vector is capable of autonomous replication inside plant cells and autonomous movement between plant cells. Such viral RNA vectors are known in the art and may be based on Potato Virus X as described e.g. in WO 93/03161, WO95/34668, Ruiz et al. (1998)

In a particularly preferred embodiment, the used viral RNA vector is derived from cowpea mosaic virus. Wellink et al. (1998) have described the use of a viral vector derived from this RNA virus for the expression of GFP in plants and demonstrated that the virus and RNA vectors derived thereof have an excellent capacity for spreading throughout an infected plant, particularly *Nicotiana benthamiana*. CPMV is an icosahedral virus with a bipartite RNA genome, consisting of a longer and a shorter RNA. Wellinck et al. have demonstrated that it is possible to incorporate extra genetic information in the shorter RNA, by inserting the GFP coding region in frame into the viral encoded polyprotein. For the purpose of the herein described methods, it is preferred that the inhibitory RNA encoding nucleic acid be inserted downstream of the polyprotein encoding open reading frame.

The inventors have obtained for the first time indications that gene-silencing may be obtained in plant cells such as protoplasts, using a viral vector derived from a satellite virus comprising a β -1,3-glucanase coding region, in a co-infection experiment with a helper virus of the satellite virus

In another preferred embodiment, the viral RNA vector is capable of replication and cell-to-cell movement, only when the required functions are provided *in trans*. Particularly preferred, is the use of a satellite virus derived RNA vector, which can replicate in plant cells and spread throughout the plant, when a corresponding helper virus is present.

As used herein, "a satellite virus" indicates an RNA virus, preferably a single stranded RNA virus, the RNA genome of which is capable of replicating in a plant cell and being encapsidated by coat protein molecules to form a virus particle or virion, only when provided externally with any number of required essential functions therefor. By "externally provided" is meant that such functions are not encoded by the satellite viral genome. Satellite viruses thus depend upon external provision of essential functions, and may lack the capacity to encode functional replicase, movement protein, or other essential functions required to complete their life cycle inside a plant cell. In a natural situation, such essential functions are usually provided by an autonomously replicating virus or so-called helper virus.

Satellite viruses useful for the present invention may include wild type isolates, but also encompassed by this definition are variants which result in reduced or minimal symptoms when infected on a host plant, particularly when co-inoculated with a corresponding helper virus. The definition also includes synthetic satellite viruses such as defective viruses and chimeric satellite viruses.

A "viral RNA vector derived from a satellite virus" should at least include cis elements from a satellite virus which are recognized by an externally provided replicase, and an origin of assembly allowing encapsidation by the provided coat protein. Preferably, the viral RNA vector does not comprise a gene encoding a functional coat protein, particularly it does not comprise the nucleotide sequence which is essentially similar to the nucleotide sequence encoding a coat protein gene. Particularly preferred are viral RNA vectors comprising an origin of assembly recognized by coat protein molecules from a rod-shaped virus, such as tobacco mosaic virus, since rod-shaped viruses do not exhibit the spatial constraints imposed on the size of genome by icosahedral viruses, thus allowing a larger number of additional nucleotides to be incorporated in the viral vector. Conveniently, the viral RNA vector comprises a number of unique or low-occurrence restriction recognition sites.

The use of viral RNA vectors derived from a satellite virus, additionally solves problems associated with the use of viral RNA vectors, such as reducing the size of the vectors, increasing versatility etc.

5 Particularly suited for the invention are viral RNA vectors derived from satellite tobacco mosaic virus comprising the origin of assembly (OAS) from TMV, preferably comprising the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5443 to the nucleotide at position 5518 or the nucleotide of SEQ ID No 5 from the nucleotide at position 5430 to the nucleotide at
10 position 5505 (such as the nucleotide sequence of SEQ ID No 12) and wherein the coat protein encoding gene has been deleted. Also particularly suited for the invention are viral RNA vectors derived from satellite necrosis vector strain comprising the OAS from TMV and wherein most of the coat protein gene has been deleted. Non-limiting examples of viral RNA vectors,
15 suitable for the invention are described hereinafter.

"A corresponding helper virus" as used herein, indicates those RNA viruses, preferably single stranded RNA viruses, which can supply the satellite virus or the derived viral RNA vector with the functions required *in trans* by that
20 satellite virus or the derived viral RNA vector, to allow it to replicate in the cytoplasm of plant cells, and spread throughout an infected plant. Typically, corresponding helper viruses will provide the satellite virus or the vector derived thereof with a replicase (RNA dependent RNA polymerase) which recognizes the *cis* sequences present on the satellite virus RNA, and will
25 allow replication of the satellite virus genome or the derived vector. Other proteins which may typically be provided by the helper virus are movement proteins, allowing inter alia, the plasmodesmata-mediated spread of viral particles from cell to cell. For satellite viruses or viral RNA vectors derived thereof which lack a functional coat protein encoding gene, corresponding
30 helper viruses may also provide a functional coat protein. Preferably, the corresponding helper virus will be capable of autonomous systemic spread in an infected plant. However, such a systemic spread seems not to be a prerequisite for efficient gene silencing. Functions required *in trans* for one

particular viral RNA vector may be supplied *in trans* by different corresponding helper viruses.

It is clear that the corresponding helper viruses may be wild type isolates of RNA viruses, preferably single-stranded RNA viruses such as the tobamoviruses or necroviruses. Particularly preferred are rod-shaped RNA viruses such as tobamoviruses including tobacco mosaic virus and the related tobamoviruses such as ribgrass mosaic virus, turnip vein clearing virus, chine rape mosaic virus, oilseed rape mosaic virus.

10

When TMV can be used as a helper virus, it can also be replaced by one of the closely related tobamoviruses mentioned above, particularly when using the viral vectors in particular plant species.

Also encompassed by the methods and means of the invention are variants of such wild type isolates, preferably variants or mutants which develop minimal symptoms when inoculated on host plants or when co-infected with a corresponding satellite virus or RNA vector derived thereof. Further preferred helper viruses may be variants or mutants of wild type isolates which have an extended host range such as tobamoviruses which can replicate and spread in corn or brassicae.

20

However, corresponding helper viruses may also be chimeric or hybrid viruses, wherein part of the viral genome has been replaced by a foreign nucleic acid, particularly wherein part of the viral genome has been replaced by a nucleic acid derived from another viral genome, preferably a part comprising a nucleotide sequence encoding a movement protein, or a part comprising a nucleotide sequence encoding a coat protein. E.g. when using a necrovirus such as TNV, it may be advantageous to insert a movement protein encoding region, preferably a movement protein derived from a tobamovirus such as TMV, particularly the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 4903 to the nucleotide at position 5709, so as to ensure spreading of the viral particles beyond the infected leaf. However,

30

spreading of the helper virus or the viral RNA vector is not essential for efficient inactivation of expression of the target genes throughout the plant, as was found by the inventors. Also when using e.g. a necrovirus such as TNV, it may be further advantageous to replace the coat protein coding region of the
5 necrovirus by a coat protein coding region of a rod-shaped virus, such as TMV, particularly the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5712 to the nucleotide at position 6191. It goes without saying that an appropriate origin of assembly for the substituted coat protein has to be incorporated in the genome of the chimaeric helper virus. In the
10 above described example however, the OAS of TMV is conveniently located within the movement protein coding region. Non-limiting examples of corresponding helper viruses will be described hereinafter.

It should be clear that whenever it is stated that plants are co-infected or
15 infected with a viral RNA vector and a corresponding helper virus, it is equal whether the helper virus is inoculated before, after or simultaneous with the viral RNA vector, provided however that there is a reasonable time limit between infection of the viral RNA vector or the corresponding helper virus.

20 Alternatively, the required functions *in trans* for the replication and movement of the viral RNA vector may be provided from the expression of chimeric genes, encoding a replicase (RNA dependent RNA polymerase) and/or a movement protein and/or a functional coat protein, integrated in the genome of the test plants.

25 Preferred kits to deliver inhibitory RNA or gene-silencing constructs to plant cells to be used in the herein disclosed methods comprise a viral RNA vector derived from a satellite RNA virus, particularly from satellite tobacco necrosis vector (STNV) or satellite tobacco mosaic virus (STMV) and a corresponding
30 helper virus, particularly a rod-shaped corresponding helper virus, wherein the viral RNA vector comprises a gene-silencing construct.

In a preferred embodiment the kit comprises a viral RNA vector derived from satellite tobacco necrosis vector, preferably comprising the cis-elements required for replication, particularly comprising the nucleotide sequence of SEQ ID No 3 from the nucleotide at position 1 to the nucleotide at position 32 and the nucleotide sequence of SEQ ID No 3 from the nucleotide at position 738 to the nucleotide at position 1245, wherein an origin of assembly of tobacco mosaic virus has been inserted, preferably comprising the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5443 to the nucleotide at position 5518 or comprising the nucleotide sequence of SEQ ID No 5 from the nucleotide at position 5430 to the nucleotide at position 5505, or comprising the nucleotide sequence of SEQ ID No 12 and wherein said helper virus is derived from tobacco necrosis virus, preferably with a nucleotide sequence of SEQ ID No 1, and comprises a gene encoding the movement protein of tobacco mosaic virus, preferably with the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 4903 to the nucleotide at position 5709 or with the nucleotide sequence of SEQ ID No 15 from the nucleotide at position 479 to the nucleotide at position 1285 and a gene encoding the coat protein of tobacco mosaic virus, preferably with the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5712 to the nucleotide at position 6191 or with the nucleotide sequence of SEQ ID No 15 from the nucleotide at position 1288 to the nucleotide at position 1767.

Preferred combinations are those kits wherein the viral RNA vector is derived from STNV-1 or STNV-2 strains (as disclosed by Ysebaert et al. 1980; Genbank Accession number M10388 or Danthinne et al., 1991 Genbank Accession M64479) and the helper virus is TNV-A (Meulewaeter et al 1990, SEQ ID No 1). Other preferred combinations are those kits wherein the viral RNA vector is derived from STNV-C (Bringloe et al. (1998); Genbank Accession Nr AJ000898) and the corresponding helper virus is TNV-D (Coutts et al. (1991); Genbank Accession Nr D00942).

In another particularly preferred embodiment the kit comprises a viral RNA vector derived from satellite tobacco mosaic virus, preferably comprising the

cis-elements required for replication, particularly comprising the nucleotide sequence of SEQ ID No 4 from the nucleotide at position 1 to the nucleotide at position 197 and the nucleotide sequence of SEQ ID No 4 from the nucleotide at position 604 to the nucleotide at position 1058 or comprising the nucleotide sequence of SEQ ID No 13 and the nucleotide sequence of SEQ ID No 14; and further comprising an origin of assembly of tobacco mosaic virus, preferably comprising the nucleotide sequence of SEQ ID No 2 from the nucleotide at position 5443 to the nucleotide at position 5518 or comprising the nucleotide sequence of SEQ ID No 5 from the nucleotide at position 5430 to the nucleotide at position 5505 or comprising the nucleotide sequence from SEQ ID No 12, and wherein said corresponding helper virus is a tobacco mosaic virus, particularly TMV-U1 (SEQ ID No 2) or TMV-U2 (SEQ ID No 5).

It will be clear to the person skilled in the art that viral RNA vectors may be generated conveniently by *in vitro* transcription methods from cDNA copies of the viral RNA. Likewise, infectious viral RNA for the corresponding helper viruses may be generated from cDNA copies of their genome. Libraries, viral vectors and corresponding helper viruses may also be maintained by replication in plant cells.

Methods to infect or inoculate plants and plant cells with viral RNA vectors, helper viruses and libraries comprised within viral RNA vectors are well within the realm of the person skilled in the art and may be performed according to the methods described in Walkey (1985).

In one embodiment of the methods of the invention, plants are inoculated, e.g. with a solution containing the libraries of gene-silencing constructs in a viral vector, or with a solution containing a mixture of gene-silencing constructs in a viral vector and corresponding helper virus. The solution may further contain additional compounds to improve inoculation and infection of the plants, such as, but not limited to abrasives, adherents, tensio-active products and the like.

Plants may be infected during different developmental stages, in order to maximize the phenotype under investigation. Also different parts of plants may be inoculated to optimize observation of the expected phenotype.

5 Although not intending to limit the scope of the invention to a particular mode of action, it is thought that the inhibitory RNA comprised within the viral RNA vector can exercise its inhibiting effect, provided there is a balance between RNA encapsidated in a virion and free RNA. It is thought that the balance between encapsidated and free RNA may be influenced by varying the
10 sequence and position of an origin of assembly within the viral RNA vector. However, the gene-silencing effect may be amplified by placing the inhibitory RNA encoding nucleic acid under control of a viral promoter, preferably a coat protein promoter, or a subgenomic promoter so that during the life cycle of the virus additional inhibitory RNA is generated or transcribed.

15

"Gene-silencing constructs" as used herein is to be interpreted as a nucleic acid, which when transcribed yield "inhibitory RNA" comprising or consisting of sense RNA or antisense RNA, or a combination of both comprising a nucleotide sequence which has at least 75%, preferably at least 80%,
20 particularly at least 85%, more particularly at least 90%, especially at least 95% sequence identity with or is identical to the nucleotide sequence whose expression is to be suppressed, or its complement. Further, the nucleotide sequence of the sense or antisense region should preferably be at least about 100 nucleotides in length, more preferably at least about 250 nucleotides,
25 particularly at least about 500 nucleotides but may extend to the full length of the coding region of the gene whose expression is to be reduced.

For practical purposes in the application of the methods for high throughput screening or validation, the gene-silencing construct may be identical in
30 sequence and length to the target nucleic acids, or they may be exactly complementary in sequence and identical in length to the target nucleic acids.

For the purpose of this invention the "sequence identity" of two related nucleotide or amino acid sequences, expressed as a percentage, refers to the number of positions in the two optimally aligned sequences which have identical residues (x100) divided by the number of positions compared. A gap, i.e. a position in an alignment where a residue is present in one sequence but not in the other is regarded as a position with non-identical residues. The alignment of the two sequences is performed by the Wilbur and Lipmann algorithm (Wilbur and Lipmann, 1983) using a window-size of 20 nucleotides or amino acids, a word length of 2 amino acids, and a gap penalty of 4. Computer-assisted analysis and interpretation of sequence data, including sequence alignment as described above, can be conveniently performed using commercially available software packages such as the programs of the Intelligenetics™ Suite (Intelligenetics Inc., CA) or the GCG Wisconsin Package.

It is clear for the person skilled in the art that the gene-silencing constructs may comprise at the same time sense and anti-sense RNA targeted towards the same nucleotide sequence whose expression is to be reduced. Preferably, the sense and antisense RNA are at least partly complementary to each other and capable of forming a stem-loop structure, since such a configuration has been shown to increase the efficiency of gene-silencing, both in occurrence and level of gene-silencing (Waterhouse et al. 1998). In the most straightforward embodiment, at least part of the target nucleic acid, preferably the complete target nucleic acid, is cloned in duplicated form, whereby the two copies are in inverted repeat, preferably separated by an unrelated spacer nucleotide sequence.

The invention also aims at providing the herein described kits in their different embodiments. It is also an object of the invention to provide the kits comprising the helper viruses and viral RNA vectors described without the gene-silencing constructs or inhibitory RNA as well as their cDNA copies, whereby the cDNA copies are under control of a promoter (which can be used in *in vitro* transcription methods available in the art) such as but not limited to

the promoters recognized by single subunit bacteriophage polymerase promoters (T7, T3, SP6 RNA polymerase specific promoters and the like).

5 The invention further relates to a method for identifying genes which are essential in plants comprising the following steps:

- 10 a) A library of random gene-silencing constructs specific for the plant is created using a viral RNA vector which is derived from a satellite RNA virus, as herein defined including all its preferred embodiments. Preferably, the library is created in a cDNA copy of the viral vector and may be generated by inserting random DNA sequences, preferably at least about 100 nucleotides in length, particularly at least about 500 nucleotides in length. The random DNA sequences may be obtained from total DNA of a plant or may represent a subset of the genome of a plant, such as DNA derived from organelles (plastids, chloroplasts, mitochondria etc.)
15 Alternatively, the library may be created by inserting cDNAs generated by reverse transcriptase from RNA, preferably mRNA obtained from said plant. The library may be normalized, e.g. as described in Takayuki et al. (1995). The library should preferably be large enough in size, i.e. contain a sufficient number of independent clones to cover the genome of the plant, according to the standards known in the art. In a preferred embodiment,
20 the library may contain duplication of the inserted nucleic acid whereby the copies are in inverted repeat. The inserted nucleic acid may be cloned downstream of a viral promoter such as, but not limited to a coat protein gene promoter or a subgenomic promoter. It will further be clear to the person skilled in the art that the relative orientation of the inserted nucleic acid, in relation to the RNA vector is only of limited importance since either
25 sense or antisense inhibitory RNA will be generated.
- b) Assay plants are infected with individual members of the library and also with a corresponding helper virus. Infection may proceed according to any
30 of the methods mentioned herein. Clearly, the DNA copy of the library should be converted into an RNA copy according to any of the methods described herein, preferably prior to the infection of the assay plants.

- 5 c) Plants developing a gene-silencing-construct-associated phenotype are identified. As used herein, a "gene-silencing-construct-associated phenotype" is meant to indicate a phenotype which is not observed when performing a mock inoculation with a viral RNA vector without gene-silencing construct, in combination with a corresponding helper virus on a similar plant. Preferred phenotypes comprise chlorosis, necrosis or any phenotype, preferably a morphological phenotype indicating that the infected tissue is inhibited or dying or deteriorating.
- 10 d) Optionally, isolating the viral RNA vector from the tissue exhibiting the gene-silencing-construct-associated phenotype according to methods available in the art for isolation of virus. Preferably, the isolated viral RNA vector comprising the gene-silencing construct of interest should be re-assayed on fresh plants to confirm the observed phenotype. The viral RNA vector can of course also be recovered from the library if the infection of
- 15 the plants was performed in an identity-preserving way.
- e) The gene silencing construct or the nucleotide sequence information thereof, may then be used to recover the corresponding genomic or cDNA clone using methods available in the art (hybridization, PCR etc.)

20 As defined herein "essential genes" of a plant, are those genes which are necessary during the normal development of a plant. As defined, essential genes may be essential for normal development only in particular developmental stages, or only in particular tissues or organs, such as e.g. flowers. Typically, inhibition of the expression of essential genes may have a

25 lethal effect on a plant or part of a plant. Preferred essential genes are those genes which result in retardation or dying of seedlings when inhibited.

It will be clear for the person skilled in the art that if the inhibition of the target nucleic acid results in a dominant effect, as is the case for inhibition of the

30 expression of essential genes, the described methods may be performed using infection of more than one viral RNA vector comprising a gene-silencing construct per plant. Care has to be taken to not dilute the phenotypic effect too much by infecting a too large number of different viral RNA vectors

comprising differing gene-silencing constructs on the same plant. It is thought that optimally any number between one and five different inhibitory RNAs may be introduced in one plant cell.

5 In yet another embodiment of the invention, a method is provided for determining the function encoded by a nucleic acid comprising a known nucleotide sequence. This nucleotide sequence may have been obtained e.g. from a genome sequencing program, including expressed sequence tags sequencing programs. In order to unravel the function of that sequence, a
10 gene-silencing construct or inhibitory RNA targeted towards said nucleotide sequence, as described in all its embodiments herein, may be introduced into a viral RNA vector derived from a satellite virus, as described herein, and used to inoculate a plant or being introduced into a plant cell, particularly into a protoplast, together with a corresponding helper virus, as described herein.

15

A large number of the embodiments described herein thus relate to a method for the introduction of inhibitory RNA in plant cells, comprising the steps of :

- a.) introducing into a plant cell, a viral RNA vector comprising inhibitory RNA or comprising a chimeric nucleic acid which when transcribed
20 yields the inhibitory RNA, wherein the viral RNA vector is derived from a satellite RNA virus; and
- b.) introducing into the same plant cell, a corresponding helper virus.

The methods of the invention can be applied to essentially all plants for which
25 viral vector and/or corresponding helper viruses are available. The methods of the invention are thought to be particularly suited for *Nicotina* spp, particularly *N. tabacum*, *N. sylvestris*, *N. benthamiana*, and other Solanacea, rice (*Oryza sativa*) corn (*Zea Mays*), Brassica spp. , cotton (*Gossypum hirsutum*), wheat, *Arabidopsis* spp., *Petunia* spp.

30

Also envisioned by the present invention are methods for developing an agronomically useful product, such as a herbicide or a transgenic plant using the herein described methods and means, further comprising the steps of

inserting a nucleic acid, involved in the determination of a particular plant trait, isolated by the methods of the invention, preferably under control of a foreign plant-expressible promoter, particularly under control of a controllable plant-expressible promoter into the genome of a plant, particularly a crop plant.

5 When essential genes have been identified according to the methods described herein, these essential genes or their encoded gene products, particularly the encoded proteins may be used in *in vitro* assays to identify compounds inhibiting the activity, particularly the enzymatic activity, which may be used as herbicides. Alternatively, a viral RNA vector encoding gene-
10 silencing constructs targeted towards essential genes may be used as herbicidal compounds.

The following non-limiting Examples describe the construction of viral RNA vectors derived from satellite viruses, and uses thereof. Unless stated
15 otherwise in the Examples, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, NY and in Volumes 1 and 2 of Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*, *Current Protocols*, USA. Standard materials
20 and methods for plant molecular work are described in *Plant Molecular Biology Labfax* (1993) by R.D.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK.

Throughout the description and Examples, reference is made to the following
25 sequences:

SEQ ID No 1: nucleotide sequence of the genome of TNV-A

SEQ ID No 2: nucleotide sequence of the genome of TMV-U1 (Genbank
Accession Nr V01408).

30 SEQ ID No 3: nucleotide sequence of the genome of STNV-2

SEQ ID No 4: nucleotide sequence of the genome of STMV (Genbank
accession Nr. M25782).

SEQ ID No 5: nucleotide sequence of the genome of TMV-U2 (Genbank Accession Nr M34077).

SEQ ID No 6: nucleotide sequence of the tomato phytoene desaturase (*pds*) encoding cDNA (Genbank Accession Nr. X59948).

5 SEQ ID No 7: nucleotide sequence of the tobacco nitrate reductase (*nia-2*) encoding cDNA (Genbank Accession Nr. X14059).

SEQ ID No 8: nucleotide sequence of the tobacco nitrite reductase (*nir-1*) encoding cDNA (Genbank Accession Nr. X66145).

10 SEQ ID No 9: nucleotide sequence of the β -1,3-glucanase(*gn-1*) encoding cDNA of *Nicotiana plumbagenifolia*.

SEQ ID No 10: nucleotide sequence of a green fluorescent protein (*gfp*) encoding region.

SEQ ID No 11: nucleotide sequence of a β -glucuronidase (*gus*) encoding region.

15 SEQ ID No 12: nucleotide sequence of an origin of assembly of a TMV-U2 strain.

SEQ ID No 13: nucleotide sequence of the leader sequence of a STMV strain

SEQ ID No 14: nucleotide sequence of the trailer sequence of a STMV strain.

20 SEQ ID No 15: nucleotide sequence of part of the genome of a TMV-U2 strain comprising movement protein and coat protein genes.

Examples

Example I : construction of the viral RNA vector kits

5 A plasmid vector for the synthesis of an infective hybrid TMV/TNV helper virus RNA is made using the following operationally linked elements:

- A T7 RNA polymerase promoter
- A nucleic acid comprising the nucleotide sequence from the nucleotides 1 to 2234 of TNV-A (nt 1 to 2234 of SEQ ID No 1), wherein the AUG codon
10 at nucleotides 2218-2220 mutated to a different codon
- A nucleic acid comprising the nucleotide sequence encoding the open reading frame for the movement protein of TMV-U1 (nt 4903-5709 of Genbank Accession Number V01408 or nt 4903-5709 of SEQ ID No 2 or nt 479-1285 of SEQ ID No 12)
- 15 • A nucleic acid comprising the nucleotide sequence from the nucleotides 2235 to 2612 of TNV-A (SEQ ID No 1)
- A nucleic acid comprising the nucleotide sequence encoding the open reading frame for the coat protein of TMV-U1 (nt 5712-6191 of Genbank Accession Number V01408 or nt 5712-6191 of SEQ ID No 2 or nt 1288-
20 1767 of SEQ ID No 15)
- A sequence comprising nucleotide 3444 to 3684 of TNV-A (nt 3444 to 3684 of SEQ ID No 1)

25 A plasmid vector for the synthesis of an infective hybrid TMV/STNV viral vector RNA is made using the following operationally linked elements:

- A T7 RNA polymerase promoter
- A nucleic acid comprising the nucleotide sequence from nucleotide 1 to 32 of STNV-2 (nt 1 to 32 of SEQ ID No 3)
- A nucleic acid comprising the origin of assembly (OAS) of TMV-U1 (nt.
30 5443-5518 of Genbank Accession Number V01408 or nt 5443-5518 of SEQ ID No 2 or nt 1018 to 1094 of SEQ ID No 15)
- A nucleic acid comprising the nucleotide sequence from nucleotide 738 to 1245 of STNV-2 (nt 738 to 1245 of SEQ ID No 3)

A plasmid vector for the synthesis of an infective hybrid TMV/STMV viral vector RNA is made using the following operationally linked elements:

- A T7 RNA polymerase promoter
- 5 • A nucleic acid comprising the nucleotide sequence from nucleotide 1 to 197 of STMV (Genbank Accession Number M25782; nt 1 to 197 of SEQ ID No 4) or the nucleotide sequence of SEQ ID No 13.
- A nucleic acid comprising the OAS of TMV-U1 (nt. 5443-5518 of Genbank Accession Number V01408; nt 5443 to 5518 of SEQ ID No 2; nt 1019-10
10 1094 of SEQ ID No 15) or of TMV-U2 (nt 5430-5505 of Genbank Accession Number M34077; nt 5430-5505 of SEQ ID No 5) such as the nucleotide sequence of SEQ ID No 12.
- A nucleic acid comprising the nucleotide sequence from nucleotide 604 to 1058 of STMV (Genbank Accession Number M25782; nt 604 to 1058 of
15 SEQ ID No 4) or comprising the nucleotide sequence of SEQ ID No 14.

Example 2. Feasibility demonstration using known endogenes or transgenes.

20 To demonstrate the feasibility of the use of the viral kits described sub example 1 for functional knockout of specific endo- or transgenes in *Nicotiana* plants, one of the following DNA fragments is inserted in the TMV/STNV or TMV/STMV hybrid vector, immediately upstream or downstream of the TMV OAS:

- 25 • a fragment of the tomato pythoene desaturase (*pds*) cDNA (comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948)
- a fragment of the tobacco nitrate reductase (*nia-2*) cDNA (comprising nucleotides 1103 to 2114 or nucleotides 5169 to 6497 of SEQ ID No 7 or
30 Genbank Accession Number X14059)
- a fragment of the tobacco nitrite reductase (*nir-1*) cDNA (comprising nucleotide 650-1212 of SEQ ID No 8 or Genbank Accession Number X66145)

- a fragment of the β -1,3-glucanase (*gn-1*) cDNA of *Nicotiana plumbaginifolia* (SEQ ID No 9 or Genbank Accession Number X07280)
- a fragment comprising a nucleotide sequence from a green fluorescent protein (*gfp*) coding region (SEQ ID No 10)
- 5 • a fragment comprising a nucleotide sequence from nucleotide 1 to 600 of the β -glucuronidase (*gus*) coding region (SEQ ID No 11)

Infective chimeric transcripts are synthesized *in vitro*, using T7 RNA polymerase with the linearized plasmid DNAs of the described vectors as
10 templates.

The TMV/STNV RNAs are mechanically inoculated on leaves of *Nicotiana benthamiana* or *Nicotiana tabacum* plants together with the TMV/TNV RNA, whereas the TMV/STMV RNAs are inoculated together with TMV-U2 virus particles or viral RNA.

15 The infected plants are scored for phenotypes, virus accumulation, and suppression of the homologous plant gene between 1 and 4 weeks after inoculation.

Plants infected with vectors containing the *pds* cDNA show a bleaching phenotype on infected leaves and silencing of the endogenous *pds* transcript.

20 Plants infected with vectors containing the *nia-2* or *nir-1* cDNA show a chlorotic phenotype on infected leaves and silencing of the endogenous *nia-2* or *nir-1* transcript, respectively.

Plants infected with vectors containing the *gn-1* cDNA show silencing of the endogenous basic -1,3-glucanase transcript.

25 Upon infection with vectors containing the *gfp* sequence, transgenic plants that normally express a *gfp* transgene show silencing of the *gfp* transgene transcript and suppression of GFP fluorescence.

Upon infection with vectors containing the *gus* sequence, transgenic plants that normally express a *gus* transgene show silencing of the *gus* transgene
30 transcript and suppression of GUS activity.

Example 3: Inactivation of phytoene desaturase in *Nicotiana benthamiana* using a TNV/STNV hybrid vector system.

5 A TNV/STNV hybrid vector system was used for the functional inactivation of a constitutively expressed endogenous plant gene . Therefore, the following STNV hybrid vectors have been constructed:

pIF9 carrying the following operationally linked elements:

- 10 • a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811;
- a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 32 of SEQ ID No 3 (STNV-2 leader);
- a nucleic acid comprising the origin of assembly (OAS) of TMV-U1 from nucleotide 5443 to 5518 of SEQ ID No 2 or Genbank Accession Number
- 15 V01408;
- a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948;
- a nucleic acid comprising the nucleotide sequence from nucleotide 806 to
- 20 1418 of SEQ ID No 3 (STNV-2 trailer).

pIF12 carrying the following operationally linked elements:

- a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811;
- 25 • a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 32 of SEQ ID No 3 (STNV-2 leader);
- a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948;
- 30 • a nucleic acid comprising the nucleotide sequence from nucleotide 742 to 1354 of SEQ ID No 3 (STNV-2 trailer).

Infective chimeric transcripts have been synthesized in vitro using T7 RNA polymerase with the linearized plasmid DNAs of the described pIF9 and pIF12 hybrid vectors as templates using standard procedures. Control in vitro transcripts have been synthesized on linearized plasmid DNAs of the precursor plasmids of pIF9 and pIF12 without the pds fragment inserts and on linearized plasmid DNA of an infective clone of the STNV wild type and of a hybrid STNV vector carrying an insert of the cat gene.

The in-vitro transcripts have been mechanically inoculated onto leaves of four weeks old *Nicotiana benthamiana* plants together with the TNV helper virus.

All infected plants were continuously scored for pds inactivation, which resulted in a phenotype showing leaf bleaching. For all inoculations necrotic lesions were observed for the inoculated leaves after 2 days post inoculation (p.i.). Within a week, necrotic lesions also occurred in upper leaves indicating the systemic spread of the viruses in *N.benthamiana*. In most infected plants, the virus symptoms have been severe but plants survived for many weeks.

Only plants, which have been infected with the hybrid vectors pIF9 and pIF12 carrying pds fragments, showed on top of the virus symptoms additional phenotypic changes. Approximately 4 weeks p.i., green upper leaves, which did not show any virus symptoms, developed bleached spots scattered all over the leaves. The bleaching was progressive and was not accompanied by necrotic lesions. Within another three weeks, the size of the spots increased constantly and the color changed from pale green to yellow to pale white-yellow. These symptoms have never been observed in plants, on which TNV/STNV wild type or TNV/STNV-deletion mutant control inoculations were carried out. Thus, this phenotype indicates the functional knockout of pds in *N. benthamiana* plants after infection with pIF9 and pIF12 hybrid viral vectors.

30

In RNA gel blot analyses performed with total RNA preparations of green and bleached spots of upper leaves of plants showing a FKO phenotype no chimeric STNV virus RNA and only very low levels of TNV helper virus RNA

could be detected. This indicates that virus induced gene silencing of pds in a specific tissue does not need to be accompanied by high levels of virus RNA.

Example 4: Inactivation of phytoene desaturase in *Petunia hybrida* using a TMV/STMV hybrid vector system.

A TMV/STMV hybrid vector system was used for the functional inactivation of a constitutively expressed endogenous plant gene. Therefore, the following STMV hybrid vectors have been constructed:

10

pVE293 carrying the following operationally linked elements:

- a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811
- a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 15 197 of STMV of SEQ ID No 4 or Genbank Accession Number M25782 (STMV leader)
- a nucleic acid comprising the origin of assembly (OAS) of TMV-U2 from nucleotide 5430 to 5505 of SEQ ID No 5 or Genbank Accession Number M34077 ("short" TMV OAS)
- 20 • a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948
- a nucleic acid comprising the nucleotide sequence from nucleotide 604 to 1058 of STMV of SEQ ID No 4 or Genbank Accession Number M25782
- 25 (STMV trailer)
- a SP6 RNA polymerase promoter comprising nucleotide 143 to 124 of Genbank Accession Number X65308

30

pVE294 carrying the following operationally linked elements:

- a T7 RNA polymerase promoter comprising nucleotide 402 to 420 of Genbank Accession Number M77811

- a nucleic acid comprising the nucleotide sequence from nucleotide 1 to 197 of STMV of SEQ ID No 4 or Genbank Accession Number M25782 (STMV leader)
- a nucleic acid comprising the origin of assembly (OAS) of TMV-U2 from nucleotide 5441 to 5849 of SEQ ID No 2 or Genbank Accession Number M34077 ("long" TMV OAS)
- a fragment of the tomato phytoene desaturase (pds) cDNA comprising nucleotide 1021 to 1671 of SEQ ID No 6 or Genbank Accession Number X59948
- a nucleic acid comprising the nucleotide sequence from nucleotide 604 to 1058 of STMV of SEQ ID No 4 or Genbank Accession Number M25782 (STMV trailer)
- a SP6 RNA polymerase promoter comprising nucleotide 143 to 124 of Genbank Accession Number X65308

Infective chimeric transcripts have been synthesized in vitro using T7 RNA polymerase with the linearized plasmid DNAs of the described pVE293 and pVE294 hybrid vectors as templates using standard procedures. Control in vitro transcripts have been synthesized on linearized plasmid DNAs of the precursor plasmids of pVE293 and pVE294 without the pds fragment insert and on linearized plasmid DNA of the infective clone STMV-10 of the wild type.

The in vitro transcripts have been mechanically inoculated onto leaves of three months old *Petunia hybrida* V26 plants together with the TMV-U2 helper virus.

The infected plants have been continuously scored for pds inactivation, which resulted in a phenotype showing leaf bleaching. The infection of petunia plants with TMV/STMV caused the occurrence of crinkled leaves starting from the infected branches but quickly progressing systemically throughout the plant. The plants did survive the infections and often showed recovery phenotypes with almost no symptoms.

Only the plants, which have been infected with the hybrid vectors pVE293 and pVE294 carrying the pds fragment, showed in addition to the virus symptoms, additional phenotypic changes. In case of TMV/pVE293 infections, the infected branches developed young leaves with bleached sectors around the veins and at the tip of the leaves. This bleaching was progressive and independent from the presence of virus symptoms. Some of the leaves were fully bleached and progressively changed into almost white color. This phenotype was restricted to leaves of the infected branches. In case of TMV/pVE294 infections a similar bleaching of young leaves was observed for all developing branches but not for branches carrying already terminal flower buds. The bleaching started again around the veins and leaf tips but proceeded quickly producing variable leaf variegation patterns. This phenotype has never been observed in control inoculations with wild type viruses or deletion mutants. Therefore this phenotype indicates the functional knockout of pds in *Petunia hybrida* plants after infection with pVE293 and pVE294 viral hybrid vectors.

In RNA gel blot analyses performed with total RNA preparations of green and bleached spots of upper leaves of plants showing a FKO phenotype no chimeric STMV virus RNA and only low levels of TMV helper virus RNA could be detected. This indicates that virus induced gene silencing of pds in a specific tissue does not need to be accompanied by high levels of virus RNA.

REFERENCES

- Ausubel *et al.* (1994) *Current Protocols in Molecular Biology*, Current Protocols, USA.
- Baulcombe (1996) *Plant Cell* 8: 1833-1844
- 5 Baulcombe *et al.* (1998) *JIC & SL Annual Report 1996/1997*
- Bringlee *et al.* (1998) 79: 1539-1546
- Chapman (1991) PhD dissertation, University of Cambridge, UK
- Coutts *et al.* (1991) *J. Gen. Virology* 72: 1521-1529
- Danthinne *et al.* (1991) *Virology* 185: 605-614
- 10 Depicker and Van Montagu (1997) *Curr. Opin. Cell. Biol.* 9: 373-382
- English *et al.* (1996) *Plant Cell* 8, 179-188
- Hamilton *et al.* (1998) *The Plant Journal* 15(6): 737-746
- Kempin *et al.* (1997) *Nature* 389: 802-803
- Kumagai *et al.* (1995) *Proc. Natl. Acad. Sci USA* 92: 1679-1683
- 15 Meulewaeter *et al.* (1990) *Virology* 177:699-709
- Pereira and Aerts (1998) *Methods in Molecular Biology* 82 Eds. Martinez-Zapatar and Salinas, Humana Press, NJ
- R.D.D. Croy (1993) *Plant Molecular Biology Labfax* jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific
- 20 Publications, UK.
- Prahasar *et al.* (1996) *Proc. Natl. Acad. Sci USA* 93: 659-663
- Ruiz *et al.* (1998) *The Plant Cell* 10: 937-946
- Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, NY
- 25 Stam *et al.* (1997) *Ann. Botan.* 79:3-12
- Takayuki *et al.* (1995) *The Plant Journal* 8(5):771-776
- Walkey (1985) *Applied Virology*, William Heinemann Ltd, London
- Waterhouse *et al.* (1998) *Proc. Natl. Acad. Sci USA* 95: 13959-13964
- Wellink *et al.* (1998) Abstract presented at the Joint Meeting of Arbeitskreis
- 30 Virologie and Nederlandse Kring voor Plantenvirologie in Wageningen, The Netherlands, November 12 and 13, 1998.
- Wilbur and Lipmann (1983) *Proc. Nat. Acad. Sci. USA* 80: 726
- Ysebaert *et al.* (1980) *J. Mol. Biol.* 143: 273-287

Claims

1. A method for isolating genes involved in the determination of a trait or phenotype of a plant species, said method comprising
 - 5 a.) Identifying a set of nucleic acid sequences of genes, whose expression is correlated with a trait of interest.
 - b.) creating a library of gene silencing constructs in a viral RNA vector, said viral RNA vector being capable of replication inside plant cells and optionally movement between plant cells of a plant, and said gene
10 silencing constructs being targeted to the nucleotide sequence of said nucleic acid sequences;
 - c.) infecting a collection of individual plants of said plant species with said library of gene silencing constructs, whereby each plant is infected with at least one member of said library;
 - 15 d.) identifying a plant wherein said trait or phenotype is altered, using an assay adapted to said trait or phenotype;
 - e.) isolating said gene involved in the determination of said trait or phenotype in said plant species, from said library based on the nucleotide sequence to which said gene silencing construct in said
20 identified plant was targeted.
2. The method of claim 1, wherein said viral RNA vector is capable of autonomous replication inside plant cells and optionally autonomous
25 movement between plant cells.
3. The method of claim 2, wherein said viral RNA vector is derived from cowpea mosaic virus.
4. The method of claim 1, wherein said viral RNA vector is capable of
30 replication inside plant cells and optionally movement between plant cells when the required factors are supplemented *in trans*.

5. The method of claim 4, wherein said viral RNA vector is derived from a satellite RNA virus and said factors are supplemented by infection with a helper virus or helper virus RNA.
- 5 6. The method of claim 5, wherein said satellite RNA virus is satellite tobacco mosaic virus.
7. The method of claim 6, wherein said viral RNA vector comprises an origin of assembly of tobacco mosaic virus.
- 10 8. The method of claim 7, wherein said helper virus is tobacco mosaic virus.
9. The method of claim 5, wherein said satellite RNA virus is satellite tobacco necrosis virus.
- 15 10. The method of claim 9, wherein said viral RNA vector comprises an origin of assembly of tobacco mosaic virus.
11. The method of claim 10, wherein said helper virus is derived from tobacco necrosis virus and comprises a coat protein gene of tobacco mosaic virus and optionally a movement protein gene of tobacco mosaic virus.
- 20 12. The method of claim 11, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said helper virus is derived from TNV-A.
- 25 13. The method of claim 11, wherein said satellite RNA virus is STNV-C and said helper virus is derived from TNV-D.
- 30 14. The method of claim 1, wherein said gene-silencing constructs comprise antisense RNA.

15. The method of claim 1, wherein said gene-silencing constructs comprise sense RNA.
16. The method of claim 1, wherein said gene-silencing constructs comprise
5 an inverted repeat.
17. The method of claim 1, wherein said gene-silencing constructs comprise complementary stretch of at least 50 nucleotides of sense and antisense RNA.
10
18. The method of claim 17, wherein said gene-silencing constructs comprise complementary stretch of at least 100 nucleotides of sense and antisense RNA.
- 15 19. The method of claim 17, wherein said gene-silencing constructs comprise at least two copies of part of the nucleotide sequences of said collection of nucleic acids, said copies being in inverted repeat.
20. A method for the isolation of a nucleic acid with a specific function from a
20 collection of nucleic acids, said collection of nucleic acids being characterized in that variation in the expression pattern of said nucleic acids is correlated with variation in a trait/phenotype of a plant harboring said nucleic acids, said method comprising the steps of
- 25 a.) creating a library of gene silencing constructs in a viral RNA vector, said viral RNA vector being capable of replication inside and optionally movement between plant cells, and said gene silencing constructs being targeted/adapted to the nucleotide sequence of said nucleic acids;
- 30 b.) infecting a collection of plants with said library of gene silencing constructs, whereby each plant is infected with at least one member of said library;
- c.) identifying plants with altered trait or phenotype using an assay adapted to said trait or phenotype.

- 5 21. The method of claim 20, further comprising the step of isolating said nucleic acid with said specific function from said identified plant with altered trait or phenotype.
- 10 22. A method for determining the function encoded by a nucleic acid comprising a known nucleotide sequence in a plant, said method comprising
- a.) providing a viral RNA vector, said viral RNA vector being derived from a satellite RNA virus, comprising a gene-silencing construct targeted to a gene comprising said known nucleotide sequence;
 - 15 b.) infecting said plant with said viral RNA vector and a corresponding helper virus;
 - c.) identifying an altered trait or phenotype of said co-infected plant.
- 20 23. A method for isolating essential genes in a plant, comprising
- a.) creating a library of random gene-silencing constructs for said plant comprised within a viral RNA vector, said viral RNA vector being derived from a satellite RNA virus;
 - b.) infecting a plant with at least one member of said library and with a corresponding helper virus;
 - 25 c.) identifying plants developing a gene-silencing-construct-associated phenotype, preferably chlorosis or necrosis.
24. The method according to claim 23, further comprising the step of isolating the viral RNA vector from the tissue exhibiting the phenotype.
- 30 25. The method of claim 23, wherein said library is created by cloning random DNA fragments of said plant in a cDNA copy of the viral RNA vector.

26. The method of claim 23, wherein said library is created by cloning random cDNA fragments of said plant in a cDNA copy of the viral RNA vector.
27. The method of claim 23, wherein said library is created by cloning random duplicated cDNA fragments in inverted repeat.
28. The method of claim 23, wherein said viral RNA vector is derived from STMV and comprises an origin of assembly of tobacco mosaic virus, and wherein said helper virus is tobacco mosaic virus.
29. The method of claim 23, wherein said viral RNA vector is derived from satellite tobacco necrosis virus and comprises an origin of assembly of tobacco mosaic virus and wherein said helper virus is derived from tobacco necrosis virus and comprises a coat protein gene of tobacco mosaic virus and optionally a movement protein gene of tobacco mosaic virus.
30. The method of claim 29, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said helper virus is derived from TNV-A.
31. The method of claim 29, wherein said satellite RNA virus is STNV-C and said helper virus is derived from TNV-D.
32. A method for the introduction of inhibitory RNA in the cytoplasm of plant cells, said method comprising:
- Introducing into said plant cell, a viral RNA vector comprising said inhibitory RNA or comprising a chimeric nucleic acid which when transcribed yields said inhibitory RNA, wherein said viral RNA vector is derived from a satellite RNA virus; and
 - Introducing a corresponding helper virus into said plant cell.


33. The method of claim 32, wherein said inhibitory RNA comprises sense RNA.
34. The method of claim 32, wherein said inhibitory RNA comprises antisense RNA.
35. The method of claim 32, wherein said inhibitory RNA comprises an inverted repeat.
36. The method of claim 32, wherein said inhibitory RNA comprises complementary stretch of at least 50 nucleotides of sense and antisense RNA.
37. The method of claim 36, wherein said inhibitory RNA comprises complementary stretch of at least 100 nucleotides of sense and antisense RNA.
38. The method of claim 32, wherein said viral RNA vector is derived from STMV and comprises an origin of assembly of tobacco mosaic virus, and wherein said helper virus is tobacco mosaic virus.
39. The method of claim 32, wherein said viral RNA vector is derived from satellite tobacco necrosis virus and comprises an origin of assembly of tobacco mosaic virus and wherein said helper virus is derived from tobacco necrosis virus and comprises a coat protein gene of tobacco mosaic virus and optionally a movement protein gene of tobacco mosaic virus.
40. The method of claim 39, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said helper virus is derived from TNV-A.

41. The method of claim 39, wherein said satellite RNA virus is STNV-C and said helper virus is derived from TNV-D.
42. The method of any one of claims 32 to 41, wherein said plant is selected
5 from *Nicotina* spp, *Oryza sativa*, *Zea Mays*, *Brassica* spp. , *Gossypum* spp., *Triticum* spp., *Arabidopsis* spp. or *Petunia* spp.
43. A kit for introduction of inhibitory RNA in the cytoplasm of a plant cell, said
10 kit comprising
a.) a viral RNA vector derived from a satellite RNA virus, said vector comprising a chimeric nucleic acid which when transcribed yields said inhibitory RNA or which comprises said inhibitory RNA; and
b.) a corresponding helper virus.
15
44. The kit of claim 43, wherein said inhibitory RNA comprises sense RNA.
45. The kit of claim 43, wherein said inhibitory RNA comprises antisense RNA.
20
46. The kit of claim 43, wherein said inhibitory RNA comprises an inverted repeat.
47. The kit of claim 43, wherein said inhibitory RNA comprises complementary
25 stretch of at least 50 nucleotides of sense and antisense RNA.
48. The kit of claim 43, wherein said inhibitory RNA comprises complementary stretch of at least 100 nucleotides of sense and antisense RNA.
- 30 49. The kit of claim 43, wherein said viral RNA vector is derived from STMV and comprises an origin of assembly of tobacco mosaic virus, and wherein said corresponding helper virus is tobacco mosaic virus.


50. The kit of claim 43, wherein said viral RNA vector is derived from satellite tobacco necrosis virus and comprises an origin of assembly of tobacco mosaic virus and wherein said corresponding helper virus is derived from tobacco necrosis virus and comprises the coat protein gene of tobacco mosaic virus and optionally the movement protein gene of tobacco mosaic virus.
51. The kit of claim 50, wherein said satellite RNA virus is satellite tobacco necrosis vector strain 1 or 2 and said corresponding helper virus is derived from TNV-A.
52. The kit of claim 50, wherein said satellite RNA virus is STNV-C and said corresponding helper virus is derived from TNV-D.

1/1

1. Tobacco mosaic virus




2. Tobacco necrosis virus



3. Satellite tobacco mosaic virus

Wild type Chimaeric



4. Satellite tobacco necrosis virus

Wild type Chimaeric



SEQUENCE LISTING

<110> Aventis CropScience N.V.

<120> Methods and means for delivering inhibitory RNA to plants and applications thereof

<130> FKOSAT

<140>

<141>

<150> US SN 09/294022

<151> 1999-04-20

<160> 15

<170> PatentIn Ver. 2.1

<210> 1

<211> 3684

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of the nucleotide sequence of the genome of TNV-A

<400> 1

```

agtattcata ccaagaatac caaatagggtg caaggcctta ctcagctaaa gagtctaaaa 60
tggagctacc aaaccaacac aagcaaacgg ccgccgaggg ttctgtatct ttcctaaact 120
ggctatgcaa cccatggaga cgacagcgaa cagtcaacgc tgcagttgcg ttccaaaaag 180
atcttctcgc cattgaggat tccgagcatt tggatgacat caatgagtgt ttcgaggagt 240
ctgctggggc acaatctcag cgaactaagg ttgtcgccga cggagcatat gcccccgcaa 300
aatccaacag gaccgcgcca gtctgtaagc agaagaagca caagtttgta aaatatcttg 360
tcaacgaagc tcgtgccgag tttggattgc ccaaaccaac tgaggcaaac agacttatgg 420
tccaacattt cttgctcaga gtgtgcaagg attggggcgt tggtactgcc cacgtacacg 480
gcaatgttgc actagctttg ccaactgggt tcatcccaac ggaagatgat ctgctatcac 540
gagcattgat gaacacacat gctactagag ccgctgtacg aggcattggac aatgtccaag 600
gggagggggt gtggaacaat aggttgggga ttgggggcca ggtcggactg gccttcgggt 660
ccaaataggg gtgccttgaa aggaggccag gattctccac gtccgtttcg cgtggggaac 720
atcctgatct ggtggtcata ccatcagggc gccctgagaa acagcgtcag ttgttacgct 780
atagtgttat aggcggccat ttattaatcg gcatccacaa caactctctt tccaacctgc 840
gtaggggctt gatggaaaga gtattctatg tcgaggggccc caatgggctt caagacgccc 900
ctaagcccggt caagggagct ttcgaaccc ttgataagtt tcgtgatctc tatactaaaa 960
atagtgtggc tcatacccct gtaactagt aacaattcct aatgaattac acgggcagga 1020
aactgactat ttacagagag gcggttgata gttgtcgca tcaaccctt agctcacgag 1080
atgcgaaact aaagacattc gtgaaggccc aaaaattaaa tctttctaag aagcctgacc 1140

```

```

ctgctccag ggtcatccaa cctagatcgc ctccgtataa cgtttgttg ggcaggtacc 1200
tccgacatta tgagcatcac gcgtttaaaa ccattgccaa gtgctttggg gaaatcacgg 1260
tcttcaaagg gtttactctg gagcaacaag gggaaatcat gcgctcgaag tggaataaat 1320
atgttaatcc cgtcgcagtc ggactcgacg ccagtcgttt cgaccaaacac gtgtctgttg 1380
aagcactcga gtatgagcat gaattttacc tcagagacta cccaaatgat aaacagctaa 1440
aatggctgct aaagcagcaa ttgtgcaacg taggaacggc attcgccagt gacggcatta 1500
taaaatacaa gaagaagggt tgtagaatga gcggagacat gaacacgagt ttgggcaact 1560
gcatttctaat gtgcgccatg gtctacgggt tgaaagaaca cttaaaccatc aatttgctcc 1620
ttgcaataaa tggggatgac tgcgtcattg tctgtgagaa agcggattta aagaaattga 1680
caagcagcat cgagccatat ttcaagcagt ttggattcaa gatggaagtg gaaaaacccg 1740
tgatatatt. tgagcgcata gaattttgcc aaacccaacc tgtgttcgat ggatcccagt 1800
acatcatggt acgcaaacct tctgtggtaa catctaaaga cgtcactagc cttatcccat 1860
gtcaaacgaa agcacaatac gcagaatggc tgcaagctgt aggtgagtgt ggcattgagca 1920
ttaacggtgg gattcctgtc atgcagaatt tctacaaaaa gctccaaact ggcattccgc 1980
gcacaaaatt caccaagacc ggcgagttcc agacgaacgg attggggtat cactctagat 2040
atatgcatag agtggcccggtgttccctcgc ctgaaacccg tttatccttc tatctagctt 2100
tcggtatcac accagacctc caagaagcat tggagatctt ctatgatacc cacaggcttg 2160
agttggatga tgttatccca actgatacct accaagtgtc aggagagcat ttgatcaatg 2220
gattacaaa ctgatgtaac ggaggacaat gtgcaaatac gcggtcgggc taggagcggt 2280
gagggtgaaga aacacaatgg ttcgggatta actggcgta agcgtcacgc ggtgagcgaa 2340
acatctcaga aatcacagca aggtactggc aatggaacta tgaccaatat agccgaagaa 2400
cagaccatta ccgtgacata caactttaac ttttaagtta tggctgcgtg tcgctgttgt 2460
gatacttcac caggtattac actattccct tactttgcaa ttctcatcct tatattggca 2520
atacttggtg tagggactcc caatcaacaa tatcaccatt ctccaagcac ttacgagtac 2580
aagactcaac acatttcgat cgcaaaatag acatggcagg aaagaagaac aacaacaacg 2640
gtcagtatat aatactgct actccagagc aacagggtgga gatagaccag cgcaacgccc 2700
gtcgtgctca aatgggtcgc atgaagaagg ctagacagcc cgttcagcga tacttacagc 2760
aacacgggtt gcgaaacgga ttgtccggta gagggggcta catagtggct cccacctccg 2820
gggggggttg cactcgaccc atagtccga aattctccaa caggggagat tccactatag 2880
tccgtaacac tgagattttg aacaaccaa tcttagcggc gctaggcgca ttcaatacaa 2940
caaactccgc actgattgca gcagcaccat catggctggc tagcatcgct gatctttaca 3000
gtaaatacag atggctctca tgtgagatca tctacattcc aaaatgcccc accactacca 3060
gtggatcaat tgccatggct ttcacatacg acagaaatga cgctgcaccc accgcaaggg 3120
ctcagctgtc acaatcttac aaggccatca attttccacc gtatgcggga tacgacggag 3180
cagcatattt gaattcgaac cagggagctg ggtcagccat cgccgttcaa cttgatgtta 3240
ccaagtggga caagccatgg taccacctta tctcctctgc cggttcggg gcgctcagcg 3300
tctcgtatca gaaccaatc tgcccccggt cccttggtg cgttagcgat gggggacccg 3360
ctactgctac tccagcaggg gaccttttca tcaagtacgt gattgagttc attgaaccaa 3420
tcaacccaac aatgaacgtc tagttctttg tactgtaact tggctaattg ctaaggtgga 3480
gtcacaccat tggagacgga gacggatcct gggaaacagg cttgacgggc ggggggtgg 3540
gccccgacg acgcatcact ccggatacca atggtacacc actatggcag ggtctgcca 3600
ggtcttgtgc accaagaacc cctggaaacg ggggggaggg gggtagcaca tatcatccag 3660
attgaggggc ctttgcccca cccc 3684

```

<210> 2

<211> 6395

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of TMV-U1

<400> 2

```

gtattttttac aacaattacc aacaacaaca aacaacaaac aacattacaa ttactattta 60
caattacaat ggcatacaca cagacagcta ccacatcagc tttgctggac actgtccgag 120
gaaacaactc cttgggtcaat gatctagcaa agcgtcgtct ttacgacaca gcggttgaag 180
agtttaacgc tcgtgaccgc aggcccaagg tgaacttttc aaaagtaata agcgaggagc 240
agacgcttat tgctaccggg gcgtatccag aattccaaat tacattttat aacacgcaaa 300
atgccgtgca ttgcgttgca ggtggattgc gatcttttaga actggaatat ctgatgatgc 360
aaattcccta cggatcattg acttatgaca taggcgggaa ttttgcacgc catctgttca 420
agggacgagc atatgtacac tgctgcatgc ccaacctgga cgttcgagac atcatgcggc 480
acgaaggcca gaaagacagt attgaactat acctttctag gctagagaga ggggggaaaa 540
cagtcctcaa cttccaaaag gaagcatttg acagatacgc agaaattcct gaagacgctg 600
tctgtcacia tactttccag acaatgcgac atcagccgat gcagcaatca ggcagagtgt 660
atgccattgc gctacacagc atatatgaca taccagccga tgagttcggg gcggcactct 720
tgaggaaaaa tgcctacacg tgctatgccg ctttccactt ctctgagaac ctgcttcttg 780
aagattcata cgtcaatttg gacgaaatca acgcgtgttt ttccgcgcgc ggagacaagt 840
tgaccttttc ttttgcacat gagagtactc ttaattattg tcatagttat tctaatatc 900
ttaagtatgt gtgcaaaaact tacttcccgg cctctaatag agaggtttac atgaaggagt 960
tttttagtcac cagagttaat acctggtttt gtaagttttc tagaatagat acttttcttt 1020
tgtacaaaagg tgtggcccat aaaagtgtag atagttagca gttttatact gcaatggaag 1080
acgcatggca ttacaaaaag actcttgcaa tgtgcaacag cgagagaatc ctcttgagg 1140
attcatcatc agtcaattac tggtttccca aaatgagggg tatggtcacg gtaccattat 1200
tcgacatttc tttggagact agtaagagga cgcgcaagga agtcttagtg tccaaggatt 1260
tcgtgtttac agtgcttaac cacattcgaa cataccagc gaaagctctt acatacgcaa 1320
atgttttgtc ctttgtcgaa tcgattcgat cgagggtaat cattaacggt gtgacagcga 1380
ggtccgaatg ggatgtggac aaatctttgt tacaatcctt gtccatgacg ttttacctgc 1440
atactaagct tgccgttcta aaggatgact tactgattag caagtttagt ctcggttcga 1500
aaacgggtgt ccagcatgtg tgggatgaga tttcgtggc gtttgggaac gcatttcctt 1560
ccgtgaaaga gaggtctctg aacaggaaac ttatcagagt ggcaggcgac gcattagaga 1620
tcagggtgcc tgatctatat gtgaccttcc acgacagatt agtgactgag tacaaggcct 1680
ctgtggacat gcctgcgctt gacattagga agaagatgga agaaacggaa gtgatgtaca 1740
atgcactttc agagttatcg gtgttaaggg agtctgacaa attcgatgtt gatgtttttt 1800
cccagatgtg ccaatctttg gaagttgacc caatgacggc agcgaagggt atagtcgcgg 1860
tcatgagcaa tgagagcggg ctgactctca catttgaacg acctactgag gcgaatgttg 1920
cgctagcttt acaggatcaa gagaaggctt cagaagggtc tttggtagtt acctcaagag 1980
aagttgaaga accgtccatg aagggttcga tggccagagg agagttacaa ttagctgggtc 2040
ttgctggaga tcatccggag tcgtcctatt ctaagaacga ggagatagag tcttttagagc 2100
agtttcatat ggcaacggca gattcggttaa ttctgaagca gatgagctcg attgtgtaca 2160
cgggtccgat taaagttcag caaatgaaaa accttatcga tagcctggta gcatcactat 2220
ctgctgcggg gtcaaatctc gtcaagatcc tcaaagatc agctgctatt gaccttgaag 2280
cccgtcaaaa gtttggagtc ttggatgttg catctaggaa gtgggttaac aaaccaacgg 2340
ccaagagtca tgcattgggt gttgttgaaa ccacgcgag gaagtatcat gtggcgcttt 2400
tggaaataga tgagcagggg gtggtgacat gcgatgattg gagaagagta gctgtcagct 2460
ctgagtcctg tgttttattc gacatggcga aactcagaac tctgcgcaga ctgcttcgaa 2520

```


acggagaacc gcatgtcagt agcgcaaagg ttgttcttgt ggacggagtt ccgggctgtg 2580
ggaaaaccaa agaaattctt tccagggtta attttgatga agatctaatt ttagtacctg 2640
ggaagcaagc cgcggaaatg atcagaagac gtgcgaattc ctcagggatt attgtggcca 2700
cgaaggacaa cgtaaaacc gttgattctt tcatgatgaa ttttgggaaa agcacacgct 2760
gtcagttcaa gaggttatc attgatgaag ggttgatgtt gcatactggt tgtgttaatt 2820
ttcttgtggc gatgtcattg tgcgaaattg catatgttta cggagacaca cagcagattc 2880
catacatcaa tagagtttca ggattcccgt accccgcccc ttttgccaaa ttggaagttg 2940
acgaggtgga gacacgcaga actactctcc gttgtccagc cgatgtcaca cattatctga 3000
acaggagata tgagggcttt gtcatgagca cttcttcggt taaaaagtct gtttcgcagg 3060
agatggtcgg cggagccgcc gtgatcaatc cgatctcaaa acccttgcat ggcaagatcc 3120
tgacttttac ccaatcggat aaagaagctc tgctttcaag agggatttca gatgttcaca 3180
ctgtgcatga agtgcaaggc gagacatact ctgatgtttc actagttagg ttaaccctta 3240
caccagtctc catcattgca ggagacagcc cacatgtttt ggtcgcattg tcaaggcaca 3300
cctgttcgct caagtactac actgttggtta tggatccttt agttagtatc attagagatc 3360
tagagaaact tagctcgtac ttgttagata tgtataaggc cgatgcagga acacaatagc 3420
aattacagat tgactcgggtg ttcaaagggt ccaatctttt tgttgacgag ccaaagactg 3480
gtgatatttc tgatatgcag ttttactatg ataagtgtct cccaggcaac agcaccatga 3540
tgaataattt tgatgtctgt accatgaggt tgactgacat ttcattgaat gtcaaagatt 3600
gcataattgga tatgtctaag tctgttgctg cgcctaagga tcaaatcaaa cactaatac 3660
ctatggtacg aacggcggca gaaatgccac gccagactgg actattggaa aatttagtgg 3720
cgatgattaa aaggaacttt aacgcacccg agttgtctgg catcattgat attgaaaata 3780
ctgcactctt agttgtagat aagttttttg atagttattt gcttaaagaa aaaagaaaac 3840
caaataaaaa tgtttctttg ttcagtagag agtctctcaa tagatgggtta gaaaagcagg 3900
aacaggtaac aataggccag ctgcagatt ttgattttgt agatttgcca gcagttgatc 3960
agtacagaca catgattaaa gcacaaccca agcaaaaatt ggacacttca atccaaacgg 4020
agtaccggc tttgcagacg atttgttacc attcaaaaaa gatcaatgca atatttgcc 4080
cgttgttttag tgagcttact aggcaattac tggacagtgt tgattcgagc agatttttgt 4140
ttttcacaag aaagacacca gcgcagattg aggatttctt cggagatctc gacagtcattg 4200
tgccgatgga tgtcttggag ctggatatat caaaatacga caaatctcag aatgaattcc 4260
actgtgcagt agaatacag atctggcgaa gattgggttt tgaagacttc ttgggagaag 4320
tttggaaca agggcataga aagaccacc tcaaggatta taccgcagg ataaaaactt 4380
gcatctggta tcaaagaaag agcggggacg tcacgcagtt cattggaaac actgtgatca 4440
ttgtgtcatg tttggcctcg atgttccga tggagaaaat aatcaaagga gccttttgcg 4500
gtgacgatag tctgtgtac tttccaaagg gttgtgagtt tccggatgtg caacactccg 4560
cgaatcttat gtggaatttt gaagcaaaac tgtttaaaaa acagtatgga tacttttgcg 4620
gaagatatgt aatacatcac gacagaggat gcattgtgta ttacgatccc cttaaagttga 4680
tctcgaaact tgggtgctaaa cacatcaagg attgggaaca cttggaggag ttcagaaggt 4740
ctctttgtga tgtgtctgtt tctgtgaaca attgtgcgta ttacacacag ttggacgacg 4800
ctgtatggga ggttcataag accgcccctc caggttcgtt tgtttataaa agtctgggtga 4860
agtatttgtc tgataaagtt ctttttagaa gtttgtttat agatggctct agttgttaaa 4920
ggaaaagtga atatcaatga gtttatcgac ctgacaaaaa tggagaagat cttaccgtcg 4980
atgtttaccc ctgtaaagag tgttatgtgt tccaaagtgt ataaataat gtttcattgag 5040
aatgagtcatt tgcagagggt gaaccttctt aaaggagtta agcttattga tagtggatag 5100
gtctgttttag ccggttttgt cgtcacgggc gagggtgaact tgcctgacaa ttgcagagga 5160
gggtgtgagcg tgtgtctggt ggacaaaagg atggaaagag ccgacgaggc cactctcgga 5220
tcttactaca cagcagctgc aaagaaaaga tttcagttca aggtcgttcc caattatgct 5280
ataaccaccc aggacgcgat gaaaaacgtc tggcaagttt tagttaatat tagaaatgtg 5340
aagatgtcag cgggtttctg tccgctttct ctggagtttg tgtcgggtgtg tattgtttat 5400

```

agaaataata taaaattagg tttgagagag aagattacaa acgtgagaga cggagggccc 5460
atggaactta cagaagaagt cgttgatgag ttcattggaag atgtccctat gtcgatcagg 5520
cttgcaaagt ttcgatctcg aaccggaaaa aagagtgatg tccgcaaagg gaaaaatagt 5580
agtaatgata ggtcagtgcc gaacaagaac tatagaaatg ttaaggattt tggaggaatg 5640
agtttttaaa agaataattt aatcgatgat gattcggagg ctactgtcgc cgaatcggat 5700
tcgttttaaa tatgtcttac agtatcacta ctccatctca gttcgtgttc ttgtcatcag 5760
cgtgggcccga cccaatagag ttaattaatt tatgtactaa tgccttagga aatcagtttc 5820
aaacacaaca agctcgaact gtcgttcaaa gacaattcag tgagggtgtg aaaccttcac 5880
cacaagtaac tgtaggttc cctgacagtg actttaaggt gtacaggtag aatgcggtat 5940
tagacccgct agtcacagca ctgtaggtg cattcgacac tagaaataga ataatagaag 6000
ttgaaaatca ggccaacccc acgactgccg aaacgttaga tgctactcgt agagtagacg 6060
acgcaacggt ggccataagg agcgcgataa ataatttaat agtagaattg atcagaggaa 6120
ccgatctta taatcggagc tctttcgaga gctcttctgg tttggtttgg acctctggtc 6180
ctgcaacttg aggtagtcaa gatgcataat aaataacgga ttgtgtccgt aatcacacgt 6240
ggtgcgtacg ataacgcata gtgtttttcc ctccacttaa atcgaagggg tgtgtcttgg 6300
atcgcgcggg tcaaatgtat atggttcata tacatccgca ggcacgtaat aaagcgaggg 6360
gttcgaatcc ccccgttacc cccggtaggg gccca 6395

```

<210> 3

<211> 1245

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of STNV-2

<400> 3

```

agtaaagaca ggaaacttta ccgactatca gaatgacaaa acgtcaaagc aaacaatcaa 60
accgcaagag cgttgcatca cagggtgcgt gtattgttga gtcaatggct gagcagaagc 120
gatttgcttt tcttacgaac accaacacag tcactacagc aggtaccgtg atcaacctga 180
gcaacaacat cgtgcaagga gatgacctg ttaatcgac cggagaccag attaagacca 240
tacaccagac tttattgact cgggtgtacag gaattaccaa cagccaaagc tttcggttca 300
tctggtttcg tgacaacacc aataggggga ctacaccggc tgtgactgag gtgtagaca 360
gtgctagtat aacatcccag tataacccca ctacgttcca gcaaaagagg ttcactgttt 420
tccaagattt catgttggt acctctatag ttggacgtgt gattgtccat cggactgccg 480
ttgataagaa acggcgtgcg atattttaca acggtgctgc ttctgtagcc gcgtcaaagt 540
gccccgggtg cacatttcta cttgtcattg gatcacatgc cactggacag tatgatgtga 600
cagccgagat tgtttatctg gacatgtaga ccattggtcat gatgatgata gtgaaggacg 660
ctgaaagatg ctagctacc ctctgggtgc acttctctgg gcaaagcaga accaaagggg 720
acgggtggtac ggccgacagt agtctgaac tagtaaata ggaccgggag aaaaccagct 780
gacggctaaa tccattccca ctagtgtatt agtggaacga ggccccgcgt gaattggggg 840
ggctgcatgg ggtggaaaac catgtggtcg cagtcatttc tctatgcat tattgtctca 900
atacttgtgt gcaacaatgc tgtaatacaa cgtagcactc aacatcactt caaaaccccc 960
tccatgtcac aagaatcaag atgcatgtct gtgtttagcg gtatatatt tgcattccact 1020
tgatcgtgat tttgccctgg gcacctcgcg cggttggtac ccgcgagac tccccacagc 1080
aacatggcat taggcagggg taaggtatag ttagtagaca aatgcgcgtg aagctggaaa 1140
gtccgggttag cagtgggggt gtgcggaatg cagcctcaac aaggtatagc tgctgcatag 1200

```

gagatgtgaa cctttcaaac ttgaattcaa gtctcatgac tgccc

1245

<210> 4

<211> 1058

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of STMV

<400> 4

```

agtaaactta ccaatcaaaa gacctaacca acaggactgt cgtgggtcatt tatgctgttg 60
ggggacatag ggggaaaaca tattgccttc ttctacaaga ggccttcagt cgccataatt 120
acttggcgcc caattttggg tttcagttgc tgtttccagc tatggggaga ggtaagggtta 180
aaccaaaaccg taaatcgacg ggtgacaatt cgaatgttgt tactatgatt agagctggaa 240
gctatcctaa ggtcaatccg actccaacgt gggtcagagc cataccttcc gaagtgtcag 300
ttcaatctgg tattgctttt aaagtaccgg tcgggtcact attttcggca aatttcggga 360
cagattcctt tacaagcgtc acagtgatga gtgtccgtgc ttggaccag ttaacaccgc 420
cagtaaatga gtacagtttt gtgaggctga agccattgtt caagactggg gactctactg 480
aggagttcga agggcgtgca tcaaacatca acacacgagc ttctgtaggg tacaggattc 540
caactaattt gcgtcagaat actgtggcag ccgacaatgt atgcgaagta agaagcaact 600
gtcgacaagt cgccttggtt atttcgtgtt gttttaactg aacctcgaca taagcctttt 660
ggatcgaagg ttaaaccgat cgctcctcgc ttgagcttga ggccggcgat ctcttatgtc 720
aacagagaca ctttgggtcta tggttgtata acaatagata gactcccgtt tgcaagatta 780
gggttaacag atcttgccgt tagtctggtt agcgcgtaac cggccttgat ttatggaata 840
gatccattgt ccaatggctt tgccaatgga acgccgacgt ggctgtataa tacgtcgttg 900
acaagtacga aatcttggtt gtgtttttcc ctccacttaa atcgaagggg tttgttttgg 960
tcttcccgaa cgcatacgtt agtgtgacta ccgttggtcg aaacaagtaa aacaggaagg 1020
gggttcgaat ccctccctaa ccgcgggtaa gcggccca 1058

```

<210> 5

<211> 6355

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
the nucleotide sequence of the genome of TMV-U2

<400> 5

```

gatgttttaa tagttttcga caacaacaat taaaacaaaa acaacatatt acaaacaaca 60
aacaacaaca atggcacaca tacaatctat aattagcaac gcccttcttg aaagcgtgag 120
tggtaaaaac actctcgta atgaccttgc aagaaggcgc atgtacgata cggccgtgga 180
agaatttaac gcccgcgacc gtagacaaa ggtcaacttt tccaaaacta ttagcgaaga 240
gcaaacgctt ctagtctcca acgctaccc ggagttccag attacctttt ataatactca 300
aaatgccgta cacagtttgg ctggaggttt gagagcatta gaattggaat atctgatgct 360
acaagttccc tatggatcgc cgacatatga tatagggtgg aactttgcag cacatttgtt 420

```

caaaggcagg gattacgtgc attgctgtat gccaatctg gacatacgag atataatgag 480
gcacgaagga caaaaggact caattgagat gtatttgtcc agattgtctc gttctaaca 540
ggtaattcct gagtttcaaa gggaggcttt taacagggtat gcagaagctc ccaacgaagt 600
ctgctgctct aaaacttttc aggattgtcg aatacatccg ccagagaata gtggtagaag 660
atacgtctgtt gctctgcaca gtttgtatga tattcctgtg catgagtttg gagctgcgtt 720
aatatctaag aatatacatg tatgttatgc agcttccatt ttggcagaag cattattact 780
agaccagacg gaggttacgc ttaatgaaat aggcgcaact ttcaaaagag aaggatga 840
tgtttctttt ttctttgctg atgaaagtac tttaaattat agtcataaat acaaaaatat 900
cttgcatat gtagttaaat ctactttcc tgcttctagt agaatagttt actttaagga 960
atttttagtc actagggtta atacttggtt ttgtaaattt accaaagtag atacctatat 1020
tctgtacaag agtgtagac aagtagggtg tgatagtga cagttctatg aggcgatgga 1080
agacgccttt gttacaaga aaaccttggc catgttcaac actgaaagag caatctttag 1140
agacacggct tcggttaact tttgggtccc taagatgaag gacatggtga tagtaccgt 1200
gtttgagggt tctattacca gcaaaaagat gacaaggagt gaggtcattg ttaatcgtga 1260
cttcgtttac acagtgtta atcatatcag aacatatcaa gccaaagcgt taacttacca 1320
gaacgtatta tctttcgttg agtctataag atcccgctg ataataatg gtgttactgc 1380
taggtctgaa tgggatgtag ataaagcaat tcttcaacc ttgtcaatga ctttcttctt 1440
gcagactaag ctggctgcgc ttcaagacga tatagtaatg ggaaagtttc ggtgcttgga 1500
taagaccact tctgaactta tttgggatga ggtgggcaaa ttttttgaa acgttttccc 1560
cactatcaaa gagagattgg tgagcaggaa aattctggat gtaagtga atgctctgaa 1620
gatcaagatc ccagatctgt atgtcacatg gaaagacagg ttcgtagctg aatacaccaa 1680
gtctgaggag ttaccgcac tagatatcaa gaaggactta gaagaagctg agcaaatgta 1740
cgacgcgtta tcagaattat ctatccttaa ggggtgctgt aatttcgata tcgcgaagt 1800
caaagacatg tgcaaggctt tagatgttag tctgtatgtg gcagcacgag taatcgttgc 1860
agtggccgag aatagaagcg gttaactct tacttttgat aagccaaccg aggagaatgt 1920
ggctaaggct cttaaaagca cggcgtctga ggccgtgga tgtcttgaa cgacatccga 1980
agaggtgaac gtaataaat tttctattgc tgagaaaggg agattgcctg tgtgtgcaga 2040
aagtcatggt ttgacgaatg ctaacttaga gcaccaggag ttggagtccc tcaacgattt 2100
ccataaggct tgcgtggata gtgtgattac aaagcaaatg gcacggttg tctacactgg 2160
ctcactcaaa gttcaacaaa tgaagaacta tgtggacagt ttggcagctt cgttgtccgc 2220
cactgtatca aatctatgca agtcaactaa ggatgaagtc gggatgatt ctgattccag 2280
ggagaaagt tgggtttggg atgtacttt gaaaaagtgg ctctcaaac ctgcggccaa 2340
aggtcattca tggggagtgt tcttgatta caaggggaaa atgtttactg cacttctatc 2400
ttatgaagga gatagaatgg tgactgagag cgactggagg aggggtggctg tatcatctga 2460
tacaatggta tattctgata ttgcaaagct ccaaaatctg agggaaaaca tgagagacgg 2520
tgaacccac gaacctactg caaagatggt acttgtggat ggggtgctg gttgtggaaa 2580
gtacaaaagga gattttgaaa gatttgatct tgataggat ttgatcttg ttctggaaa 2640
acaagctgct gctatgatca gaagaagggc taattcatct ggactgataa gagccacaat 2700
ggacaatgtg agaacggtag attcacttct aatgcatcca aaaccgcat cacacaagag 2760
gctttttatt gatgaagggt tgatgctgca caccggttgt gttacttcc tgggtgcttat 2820
ctctggttgc gacatcgcat acatttacgg agatacacag cagattcctt tcattaacag 2880
agttcagaat tccccgtatc ccaaacattt tgagaagctg caagtggatg aagttgagat 2940
gaggaggacc aactgagat gccagggtga tgtgaatttt tcttacaat cgaagtacga 3000
aggagcgggtg acaaccactt caactgtaca acgatcggc tcatctgaga tgataggcgg 3060
taaggagta ctaaacagt tttccaaacc actaaaaggg aaaattgtaa ctttactca 3120
ggctgataaa tttgagttag aggagaagggt ctataagaat gtgaacaccg ttcattgagat 3180
ccaaggagaa acctttgaag atgtgtcgct ggtagattg acggcaactc cactgactct 3240
gatttccaag tcttccccgc atgttctagt cgctctgact agacacacaa agagcttcaa 3300

atattacacc gtagtgtag atccttttagt acagataatt agtgatttgt cttctttaag 3360
 ctcttcctt ttagaaatgt atatggtaga agcaggtagt agatagcaat tacagatgga 3420
 tgcagtgttc aaaggtcata atctctttgt ggcaacacct aaatcaggag actttccaga 3480
 tctacagtgc tattacgatg tatgcctccc tggtaatagt actatactta acaagtatga 3540
 tgctgttacc atgagggttac gtgataatag tcttaatgtg aaggattgtg ttcttgattt 3600
 ttccaaaagt attccgatgc caaaggagggt gaaaccatgt ctagagccag ttttgctac 3660
 cgcggcggaa ccgccaaggg ctgcaggact actcgaaaat ctggttgcaa tgattaaaag 3720
 aaatttcaac gcaccagacc tgacggggac gattgacatt gagagcaccg catctgttgt 3780
 agtagataag ttttttgata gctattttat taaaaaagaa aaatacacaa aaaatattgc 3840
 tggagtgtg acgaaggatt caatgatgag atggttgga aacaggaaag aagtactatt 3900
 ggacgacttg gctaactaca attttacaga tctgccggcc atcgatcagt acaagcacat 3960
 gatcaaggct caacccaaac agaaattgga cctttcaatt cagaatgaat accctgctct 4020
 gcaaacaatt gtctaccatt cgaagcagat caacggtatt ttggccggtt tctcagagct 4080
 tacaagggtg ctgctcgagg catttgattc taagaagttt ctttcttta ctaggaaaac 4140
 tccagaacag attcaagaat ttttctcgga tctcgactcg cacgttccta tggatgtgtt 4200
 agaactggat atttctaagt atgataagtc acagaacgag ttctattgtg ctgtagagta 4260
 tgaaatatgg aaaagattgg gtctcaatga gtttttggcc gaagtgtgga aacaagggca 4320
 caggaaaaca actttgaagg attacattgc tggaaatcaag acatgtctgt ggtatcaaag 4380
 gaaaagcggg gatgtgacta ctttcatcgg caatactgtt ataatagcag cttgcttggg 4440
 ttcaatgta ccgatggaaa aggtcataaa aggtgctttt tgtggagacg attccgtttt 4500
 gtattttcca aagggttttg atttccctga cattcagtc tgtgctaate tcatgtggaa 4560
 ttttgaggcc aaactgtata gaaagaggta cggttacttt tgtggtagat acatcataca 4620
 ccatgataag ggagcaatag tgtattatga tcctttgaag ttgactcca aacttggggc 4680
 aaaacatatc aaggattatg atcacttaga agagttaagg gtgtctttgt gcgatgttgc 4740
 ttgttcgctc ggaaactggg gcttaggctt tccgcagctg aacgcagcta tcaaggagg 4800
 tcataaaacc gcgattgatg gttcgtttgc ttttaattgt gttacaaat ttttgtgtga 4860
 taaattttta tttagaactt tgtttttaa tggctgttag tctcagagat actgtcaaaa 4920
 ttagcgagtt cattgatctt tcgaaacagg atgagatact tccggcattc atgactaagg 4980
 tcaagagtgt tagaatatcg actgtggaca agattatggc tgttaagaat gatagtcttt 5040
 ctgatgtaga ttactttaaa ggtgttaagt tagttaagaa agggatatgtg tgcttagctg 5100
 atttggtagt gtctggggag tggaaatctc cggataactg ccgtgggtggg gtcagtgttt 5160
 gtattgtaga taagagaatg aaaaggagta aggaagcaac gctgggtgcg tatcacgccc 5220
 ctgcttgcaa aaagaatttt tcttttaagc taatccctaa ttattcaata acatccgagg 5280
 atgctgagaa gcaccctgg caagtgttag tgaatatcaa aggagtggct atggaagaag 5340
 gatactgtcc tttatctttg gagttcgttt caatttgtgt agtacataaa aataatgtaa 5400
 gaaaagggtt gagggaaactg attttgagtg tgacagacgg ctgcaccaatt gaactcactg 5460
 aaaagggtgt tgaggagttc gtggatgaag taccaatggc tgtgaaactc gaaaagggtt 5520
 cggaanaaaa aaaagaaatg gtaggtaata atgttaataa taagaaaata aataacagt 5580
 gtaagaaggg ttttaaaatt gaggaaattg aggataatgt aagtgatgac gagtctatcg 5640
 cgtcatcgag tacgttttaa tcaatatgcc ttatacaatc aactctccga gccaatgtgt 5700
 ttacttatct tccgcttacg cagatcctgt gcagctgac aatctgtgta caaatgcatt 5760
 gggtaaccag tttcaaacgc aacaagctag gacaacagtc caacagcaat ttgcggatgc 5820
 ctggaaacct gtgcctagta tgacagttag atttccctga tccgatttct atgtgtatag 5880
 atataattcg acgcttgatc cgttgatcac ggcgttatta aatagctttg atactagaaa 5940
 tagaataata gaggttgata atcaaccgc accgaatact actgaaatcg ttaacgcgac 6000
 tcagagggta gacgatgcta ctgtagctat aagggttca atcaataatt tggctaata 6060
 actggttcgt ggaactggca tgttcaatca agcaggcttt gagactgcta gtggacttgt 6120
 ctggaccaca actccggcta cttagctatt gctgtgagat ttcctaaaat aaagtcgctg 6180

aagacttaaa attcaggggtg gctgatacca aaatcagcag tgggtgttcg tccacttaaa 6240
 tataacgatt gtcatatctg gatccaacag ttaaaccatg tgatgggtga tactgtggta 6300
 tggcgtaaaa catcggagag gttcgaatcc tcccctaacc gccggtagcg gccca 6355

<210> 6

<211> 2346

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequenc : nucleotide
 sequence of the tomato phytoene desaturase (pds)
 encoding cDNA

<400> 6

cttttactag ttatagcatt cggatctttt ttctgggtaa ctgccaaacc accacaaatt 60
 acaagtttcc atttaactct tcaacttcaa cccaaccaa tttatttcct taattgtgca 120
 gaaccactcc ctatatcttc taggtgcttt cattcgttcc gaggtaagaa aagatttttg 180
 tttctttgaa tgctttatgc cactcgttta acttctgagg tttgtggatc ttttaggcga 240
 cttttttttt ttttgtatgt aaaatttggt tcataaatgc ttctcaacat aaatcttgac 300
 aaagagaagg aattttacca agtatttagg ttcagaaatg gataattttc ttactgtgaa 360
 atatccttat ggcaggtttt actgttattt ttcagtaaaa tgcctcaaat tggacttggt 420
 tctgctgtta acttgagagt ccaaggtagt tcagcttate tttggagctc gaggtcgtct 480
 tctttgggaa ctgaaagtcg agatgggtgc ttgcaaagga attcgttatg ttttgcgtgt 540
 agcgaatcaa tgggtcataa gttaaagatt cgtactcccc atgccacgac cagaagattg 600
 gttaaggact tggggccttt aaaggctcgt tgcattgatt atccaagacc agagctggac 660
 aatacagtta actatttgga ggctgcattt ttatcatcaa cgttccgtgc ttctccgcgc 720
 ccaactaaac cattggagat tgttattgct ggtgcaggtt tgggtggttt gtctacagca 780
 aaatatttgg cagatgctgg tcacaaaccg atactgctgg aggcaaggga tgttctaggt 840
 ggaaaggtag ctgcatggaa agatgatgat ggagattggt acgagactgg tttgcatata 900
 ttctttgggg cttacccaaa tattcagaac ctgtttggag aattagggat taacgatcga 960
 ttgcaatgga aggaacattc aatgatattt gcaatgccaa gcaagccagg agaattcagc 1020
 cgctttgatt tctccgaagc tttaaccgct cctttaaatt gaatttttag catcttaaag 1080
 aataacgaaa tgcttacatg gccagagaaa gtcaaatttg caattggact cttgccagca 1140
 atgcttgagg ggcaatctta tgttgaagct caagatggga taagtgttaa ggactggatg 1200
 agaaagcaag gtgtgccgga cagggtgaca gatgaggtgt tcattgctat gtcaaaggca 1260
 ctcaacttta taaaccctga cgaactttca atgcagtga ttttgatcgc attgaacagg 1320
 tttcttcagg agaaacatgg ttcaaaaatg gccttttttag atggtaatcc tcttgagaga 1380
 ctttgcattg cgattgttga acacattgag tcaaaagggt gccaaagtcag actgaactca 1440
 cgaataaaaa agattgagct gaatgaggat ggaagtgtca agagttttat actgagtga 1500
 ggtagtgcaa tcgagggaga tgcttttgtg tttgccgctc cagtggatat tttcaagctt 1560
 ctattgcctg aagactggaa agagattcca tatttccaaa agttggagaa gttagtcgga 1620
 gtacctgtga taaatgtaca tatatggttt gacagaaaac tgaagaacac atatgatcat 1680
 ttgctcttca gcagaagctc actgctcagt gtgtatgctg acatgtctgt tacatgtaag 1740
 gaatattaca accccaatca gtctatgttg gaattgggtt ttgcacctgc agaagagtgg 1800
 atatctcgca gcgactcaga aattattgat gcaacgatga aggaactagc aacgcttttt 1860
 cctgatgaaa tttcagcaga tcaaagcaaa gcaaaaatat tgaagtacca tgttgtcaaa 1920
 actccgaggt ctgtttataa aactgtgcca ggttgtgaac cctgtcggcc tttacaaaga 1980

tccccaatag aggggtttta tttagccggt gactacacga aacagaaata cttggcttca 2040
 atggaaggcg ctgtcttatac aggaagctt tgtgctcaag ctattgtaca ggattatgag 2100
 ttacttggtg gacgtagcca aaagaagttg tcggaagcaa gcgtagttta gctttgtggt 2160
 tattatttag cttctgtaca ctaaatttat gatgcaagaa gcgttgtaga caacatatag 2220
 aagaagagtg cgaggtgaag caagtaggag aaatgtagg aaagctccta taaaaagga 2280
 tggcatgttg aagattagca tctttttaat cccaagtta aatataaagc atattttatg 2340
 gaattc 2346

<210> 7

<211> 7096

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: nucleotide
 sequence of the tobacco nitrate reductase (nia-2)
 encoding cDNA

<400> 7

tacatacaag ggcggaata aacttttttt aaagtaaag tatatgaact tgcaatgaaa 60
 gaggacctta acttgtttgt ctttggtgct ttctgcaaat ttcaccttaa cagcccattt 120
 gagattgatt tagttagtta taacaattag ttaaatgctt gtgtaatttg aagaaaatat 180
 ttggacgtgc tcgctgaaaa cattatactc ctatataata gaaatacttt ctgaaaagtt 240
 ggtcttggtc aaaaacgtat aagagagttg gtcttctcat aaatagtcac tagctttctg 300
 attttttttc actttctata tcacgtaaat aggtaactcaa atttgatatt tacaccaaac 360
 aaatgaaaat aggatattgtg tttttcatat gtatatattt ctatcgtact taatgatata 420
 tacatataca tataacctta ctttttgatt actaaaaatt taattatatt taatttgggt 480
 aaatatcaga tgccacaaaa catttaccta gccactggtt ttgactacta aaaatttaat 540
 tatgttttagc ttgggtaaat atcagatgtc actaaacatt ttacctagcc attcctccga 600
 aaagaaattg agaaggaaat tagagttagt ggagccataa taatgtttta tgtgaccata 660
 actcggtgaa aaccacggca agaataagaa acagctgtta aggctaacca acagctgcat 720
 atctttaagc catttgctat taccccaaca tcgcatcttc ctctgatecc gaccctacgg 780
 gcgtaaaaag tgtaaatcgt tagaattggt ttatttattt tatgatgtca ctatttttta 840
 aaatcaaaat taaattgggg tgcgattttt tttgggtcct gcttatgtat agtatggcgc 900
 tatggaggca ctgagagagt ccgaaacgtt tctatataag gccaccccac gcattcacaa 960
 acttcgttcc caaacagaac aagaaaatca aatctcggag agagagagag agaaatattt 1020
 tgagagagaa atacagaaaa tctctcttcc tctctctctt ttttttcaa tccccattca 1080
 tattcttttt ttagaataat ctatggcggc atctgtcgaa aacaggcagt tcagtcacct 1140
 agaagccggt ttatcccggg ctttcaagcc ccggtctgat tccccggttc gtggctgcaa 1200
 cttcccttcg cccaacagta ctaatttcca aaagaaacca aattccacca tttaccttga 1260
 ttactcgtcg agtgaagacg acgatgatga tgacgaaaae aatgagtacc ttcaaatgat 1320
 taaaaaaggg aattcagagt tagagccatc tgrtcatgac actagggacg aaggtaccgc 1380
 tgataattgg attgaacgca acttttccat gattcgtctc accggaaagc atccatttaa 1440
 ctccgaacca ccgttgaaac ggctcatgca ccacggcttt atcacaccgg tcccaattca 1500
 ttacgttcgt aaccatggac cggttcccaa gggcacgtgg gatgactgga ccgtggaagt 1560
 cacgggacta gtgaagcgtc ctatgaaatt cacaatggac cagttgggta acgaattccc 1620
 ttgtagagaa ttgcccggtt cgcttggttg tgctggcaat cgaaggaaag aacagaacat 1680
 ggttaaacia accattgggt tcaactgggg ccgcgctgcc gtttcaacia cgatatggcg 1740

cggggtaccc ctccgcgctt tgctaaaacg gtgcggtggt tttagcaaga ataaaggggc 1800
gcttaatggt tgcttcgaag gagctgatgt gttgcccgga ggtggtggtt caaagtatgg 1860
aaccagcatt aagaaggaat ttgcaatgga tccagcacga gatatcatcg tagcctacat 1920
gcagaacgga gaaaaattgg cccccgacca cgggtttcca gtacgaatga taattccagg 1980
attcattgga ggaagaatgg tgaaatggat aaagaggatt atagtcacca cccaagaatc 2040
agacagctat tatcatttca aggacaatag agttcttctt ccccatgttg atgtgaact 2100
tgcaaatacc gaaggtagct accgtaacta tttcaattta ttactccatt tgttccaatt 2160
tatgtgaacc tatttctctt ttggtccggt caaaaaagaa tgaacccttt ctaaatttgg 2220
taacaattta gcttaactt acaacttcac ccttaatgag aaacttttat aaccacacaa 2280
ataccctggg gccatttgg acttgtttag gtcgacaaat tccaaaagtt ttattttttt 2340
cttaaaacttc gtgctcagtc aaacaggttc acgtaaattg aaacggagag agtatcattt 2400
ttattaaggg gtataaatat attttaatta gttgagactt gcacatacaa gtaaaatatt 2460
tcttagaata caaaatcaac tgaaagctta cttctaatta tatggttttg aattttctct 2520
tcaatgaagt aaataaaaag gaaacaatta tattcaacgc atgtaggtat atggtcctgt 2580
cattatctca aatcaaatgg tttaaagaca aaggactttg gaaacataga attgtcagct 2640
ttatagttat ggagtactat attagtttagc tgtttgcac tttcataat tggctctatct 2700
gtgtgcagca tgggtgtaca agccagagta tatcatcaat gagcttaata ttaactctgt 2760
cattacgacg ccgtgtcatg aagaaatttt gccaatatg gcctggacga ctcagcgacc 2820
ttacacgttg aggggctatt cttattcttg ttagtatttt tatattttcc gattttgctg 2880
agaatatcat atttcttagt tttgtcgata catcgatcc tctaactctg acgttttact 2940
tcgtccttat gcaccactt acgtccttac tttctcagac agtttattga tgaaaactac 3000
ttactatttt cgaccgata gcctcagcgt ccttaattaa atgtgatgtt ttgaaagaga 3060
tattctctcc cgtctatttt aattaatttt tggctgtttt tatacgtggg aatctatttt 3120
taacattaat taatatagaa atgaaccata ttaattattat taatttcttc attgaaaata 3180
caacaaatac tcttcggctc ttactacaat gacaattttg aagaaaaata attaattcct 3240
tcctaataac tgaaaaatca aatattgttg accataaaaa aagggtcaaaa aattaattaa 3300
aatgaactgg agagagtaaa ttagaaaaata taattatagc actagtaatt aaagtatta 3360
gatgtcttct ttaaaaagcg tgtgaaaact ttaagacga aatataatat gaattattatc 3420
taatacttag aaagtgtcaa taattggtag acaattttaa ctatatacta gttaaaaagt 3480
ctgtcaatac aactattagt attggggatt agagagaata gtagtaaaat ggagtaattg 3540
gacgcagtag cttgggcatg ctgattgctg tcagcttgtt tgctaattgtg aaaaagaaaa 3600
tagtaagaaa aggccaacat ggttttgttt attttattat gtggtagtac acaaaaacct 3660
ggggagcttt cctagttctg aagagtcggt ctttggttagc acaaaattaa tagtatagta 3720
taccaagtga atattaaatt caattgtcta aagcacggaa tctttttgac tacttttagt 3780
cctgcattctt gggttgcctc aacaacaccc tttattgaat tattatagta atgttcaata 3840
taatatacaa ttagaaaaca ctctaagtgg tcactttata tggatctagt caatactatt 3900
tcttctaacc aacgtgccta attacttccc actttccagt acatgaccac cattaagttt 3960
aatttttgtc aattccttgt gcaattggcc cttcaaatga gcagaagtgt tacgtaggaa 4020
aactaacttc agctactatt ataggagtaa acctgttagg aaaagatgct cgaggaactg 4080
acaaaacttg tagaataatt agccattgta ttgattgaaa tactgattgt gaacgtgtaa 4140
caaacaggcg gagggaaaaa agtaacgcga gtagaagtga cgttggatgg aggagaaaca 4200
tggaagttta gcacactaga tcaccagag aagcccacca aatatggcaa gtactggtgt 4260
tgggtctttt ggtcactcga ggttgagggt ttagacttgc tcagtgtctaa agaaattgct 4320
gttcgagctt gggatgagac cctcaatact caaccgaga agcttatttg gaacgtcatg 4380
gtacgttcac ttcttctttt acctttattt cttttaactt ctatatacta gcggtgtaaa 4440
gttattttac accataagtt aacttacaaa aatatgtaac tatctatact acgagtgtatg 4500
agggcaagaa ggggtttaag tatttgacaa taaatgtaaa ccctgcaatt ttgttcttaa 4560
ttttttatcc tttcaactct ttgtgattgc ttcattatct agattcacag agcacatgtg 4620


```

ttcacatgcc aaaacaaaaa actacaaaca aaaaaacttt tcaactagctt tagtctaaga 4680
ttccccctttt tttttttggg aggtgtgtgg tccatactcc atagatcaat tccagccact 4740
gacgtaccaa accctgaaaa ttcctagtag ttatagcgac gtacaatcat ttcattattat 4800
gtaagcagag acgtgatcac atgaactaga tgtgaatacc acttgcccag tccaccaggt 4860
caattcatct agatgtgtaa atcttgacac cagcactggg tcaacttttat aacactagca 4920
tttaacaaca tttcatcctt gaacattact tgggctaatt aataagtatt tttttttata 4980
tactctaaaa attgtaatta cataaatgaa tttaacttat acacgctgac aatgttacta 5040
attccacttt ttacggacgg ttatctatag aaatcattta ggtgaaacaa ttctcttaca 5100
ctatgatcag tgttagtaca taatggttat tacattttct aaatattgtg ctatgttgca 5160
atgttcaggg aatgatgaat aattgctggg tccgagtaaa gatgaatgtg tgcaagcctc 5220
acaagggaga gattggaata gtgtttgagc atccgactca acctggaaac caatcaggtg 5280
gatggatggc gaaggagaga catttggaata tatcagcaga ggcacctcaa aactaaaga 5340
agagtatctc aactccattc atgaacacag cttccaagat gtactccatg tccgaggtca 5400
ggaaacacag ctctgctgac tctgcttggg tcatagtcca tggcatatc tatgacgcca 5460
cgcgtttctt gaaagatcac cctgggtggg ctgacagcat tctcatcaat gctggcactg 5520
attgcaactga ggaatttgat gcaattcatt ctgataaggc taagaagctc ttggaggatt 5580
tcaggattgg tgaactcata actactgggt acacctctga ctctcctggc aactccgtgc 5640
acggatcttc ttccttcagc agctttctag cacctattaa ggaacttgtt ccagcgaga 5700
ggagtgtggc cctaattcca agagagaaaa tcccatgcaa actcatcgac aagcaatcca 5760
tctcccatga tgtaggaaa tttcgatttg cattgccctc tgaggatcaa gtcttgggct 5820
tgctgtttgg aaaacatata ttcctctgtg ccgttattga cgataagctc tgcatgcgcg 5880
cttacacgcc tactagcacg atcgatgagg tggggactct cgagttgggt gtcaagatat 5940
acttcaaagg aattcacctt aaattcccca atggagggca aatgtcacag tatcttgatt 6000
ctatgccgtt agggtcattt ctcgacgtga aagggtccatt aggtcacatt gaataccaag 6060
gaaagggaaa tttcttagtt catggcaaac agaagtttgc caagaagttg gccatgatag 6120
caggtggaac aggaataact ccagtgtatc aagtcatgca ggcaattctg aaagatccag 6180
aagatgacac agaaatgtat gtggtgtatg ctaacagAAC agaggatgat attttactta 6240
aggaagagct tgattcatgg gctgagaaaa ttccagagag ggttaaagtt tggatatgtg 6300
ttcaggattc tattaaagaa ggatggaagt acagcattgg ttttattaca gaagccattt 6360
tgagagaaca tatccctgag ccattctaca caaactggc tttggcttgt ggaccacctc 6420
ctatgattca atttgctgtt aatccaaact tggagaagat gggctatgac attaaggatt 6480
ccttattgggt gttctaattt taaaaacaaa acaatatctg caggaataaaa tttttttttt 6540
ccccctatca gttgtacata ttgtatttgg tttatcacc ccatgtacta cgtagtggtt 6600
gtagttctta catttttatt ttttagaatt tttttaaacc ttaggatata aagggtttct 6660
cttccaacaa agtgattctt tagggaagaa atgtactgta ctgtactagt atgtctaagc 6720
cgaaagttgt aatgtttacc atgacaaatt gtattcaatt cctcatggaa tagtaacatt 6780
gtgttcatgt gtcttcctgt aagcgatctt caaaatatca atgtatatat atagtaattg 6840
caaaccattg ttccttttcc cgatgtagtt aactactctt tctttagctt ctagtctctg 6900
gtgaatatatt ttttttctat aactctttta ttaatacggc cttaaataag agaaaagttt 6960
aaaccacgaa tatcattatg cagacgtata ggtaattaat ctactttttg aaaaaaaatc 7020
tattttcttt atgtgttcct tcaaaataat attctagaac cttttgtata ttccctttta 7080
acttctattt agtttt

```

7096

<210> 8

<211> 1839

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: nucleotide
sequence of the tobacco nitrite reductase (nir-1)
encoding cDNA

<400> 8

```

tttctattaa atttctggca ccttcattgc caaatccagc tagattttcc aagaatgctg 60
tcaagctcca cgcaactccg ccgtctgtgg cagcgccgcc agctggtgct ccagaggttg 120
ctgctgagag gctagaaccc agagttaggg aaaaagatgg ttattggata ctcaaggagc 180
agtttagaaa aggcataaat cctcaagaaa aggtcaagat tgagaagcaa cctatgaagt 240
tgttcatgga aaatggtatt gaagagcttg ctaagatacc cattgaagag atagatcagt 300
ccaagcttac taaggatgat attgatgtta ggcttaagtg gcttggcctc ttccatagga 360
gaaagaacca atatgggagg ttcattgatg gattgaagct tccaaatgga gtaacaacga 420
gtgcacagac tcgatacttg gcgagtgtga taaggaaata cgggaaagaa ggatgtgctg 480
atattacaac gagggcaaat tggcagattc gtggagttgt actgcctgat gtgcccga 540
tactaaaggg actagcagaa gttgggttga ccagtttgca gagtggcatg gacaatgtca 600
ggaatccagt aggaatccct cttgctggaa ttgatccaga agaaatagta gacacagggc 660
cttactactaa tttgctctcc caatttatca ctggcaattc acgaggcaat cccgcagttt 720
ctaacttgcc aaggaaagtg aatccgtgcg tagtaggctc tcatgatctt tatgaacatc 780
cccataatcaa cgatctcgcg tacatgcctg ccacgaaaga tggacgattt ggattcaacc 840
tgcttggtggg tgggttcttc agcgcaaaaa gatgtgatga ggcaattcct cttgatgcat 900
gggttccagc tgatgatgtt gttccggttt gcaagcaat actggaagct tttagagatc 960
ttggtttcag agggaaacaga cagaaatgta gaatgatgtg gtaaatcgat gaactgggtg 1020
tagaaggatt cagggcagag gtcgagaaga gaatgccaca gcaagagcta gagagagcat 1080
ctccagagga cttggttcag aaacaatggg aaagaagaga ttatcttggg gtacatccac 1140
aaaaacaaga aggctacagc tttattgggtc ttcacattcc agtgggtcgt gttcaagcag 1200
acgatatgga tgagctagct cgtttagctg atgagtatgg ttcaggagag atccggctta 1260
ctgtggaaca aacattatt attcccaaca ttgagaactc aaagattgag gactgctca 1320
aagagcctgt tctgagcaca ttttcacctg atccacctat tctcatgaaa ggtttagtgg 1380
cttgtagctg taaccagttt tgtggacaag ccataatcga gactaaagct cgttcctga 1440
tgataactga agaggttcaa cggcaagttt ctttgacacg gccagtggag atgcactgga 1500
caggtgccc gaatacgtgt gcacaagttc aagttgcgga cattggattc atgggatgcc 1560
tgactagaga taagaatgga aagactgtgg aaggcgccga tgttttctta ggaggcagaa 1620
tagggagtga ttcacatttg ggagaagtat ataagaaggc tgttccttgt gatgatttg 1680
taccacttgt tgtggactta ctagttaaca actttggtgc agttccacga gaaagagaag 1740
aaacagaaga ctaataaaat ttagaatagt tgggtgattt gctgtgttca taacatgtaa 1800
tgtatgataa atcaatgcaa acatttctac ctacgtgag 1839

```

<210> 9

<211> 1294

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA of the
beta-1,3-glucanase of *Nicotiana glauca*

<400> 9

```

ttgctcttca aatggctgct attatactgc taggattgct tgtttccagc actgagatag 60
taggagctca atcagtaggt gtttgctacg gaatgctggg caacaacttg ccaccagcat 120
cacaagttgt acaactgtac aagtcaaaaa acataagaag aatgaggctt tatgatccaa 180
atcaagcagc tttacaggct ttaagaggct ccaacattga agttatgtta ggagttccca 240
attcagatct ccaaaacatt gctgctaacc cctcaaatgc aaataattgg gtccagagga 300
atgtcagaaa tttctggcca gccgttaa attaggtacat tgccgttgga aatgaagtca 360
gccctgtaac aggacatct tcaactaccc gatattctt tccggccatg aggaacattc 420
ggaatgcgat ttcttcagca ggtttgcaa acaatatcaa agtctcaagt tctgtagaca 480
tgaccttgat tgggaactct tttccaccat cacagggttc gtttaggaac gacgttaggt 540
cgttcattga tccgattatt ggggttgtaa ggcgcataaa ttcgccttta ctcgttaaca 600
tttatcctta ttttagctat gctggtaatc cgcgcgatat ttctctcccc tatgtctttt 660
tcaactgctcc aaatgtggtg gtacaagatg gtacacttgg atatagaaac ttatttgatg 720
caatgtcgga tgctgtgtat gctgccctgt ctcgagccgg agggggctcg atagagattg 780
ttgtgtccga gagtggctgg ccatctgctg gcgcatttgc cgcgacaaca aacaatgcag 840
caacttacta caagaactta attcagcatg ttaaaagggg tagtccaaga aggcctaata 900
aagtcattga gacctattta tttgctatgt ttgatgagaa taacaaaaac cctgaattgg 960
agaaacattt tggactcttt tcccccaaca agcagccc aaatccactc agctttgggt 1020
tttcagatag atattgggac atttctgctg aaaataatgc tactgcagct tctctcataa 1080
gtgagatgtg ataagagagt tctctttaa tatctttaca tggatggaaa acttagtacc 1140
aataactaga ttgtttcttt ctttatgcaa tttcttgta atgagagact agtacttgct 1200
ctctgtgtcc ttgtggagag taactagaga caaattaagc aaataacata aataattgag 1260
tgttgattct gcaatgataa atagaaaaaa aaaa 1294

```

<210> 10

<211> 720

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: green
fluorescent protein encoding region

<400> 10

```

atggtgagca agggcgagga gctgttcacc ggggtggtgc ccatcctggt cgagctggac 60
ggcgacgtaa acggccacaa gttagcgtg tccggcgagg gcgagggcga tgccacctac 120
ggcaagctga ccctgaagtt catctgcacc accggcaagc tgcccgtgcc ctgggccacc 180
ctcgtgacca ccctgacctt cggcgtgcag tgcttcagcc gctaccccca ccacatgaag 240
cagcacgact tcttcaagtc cgccatgccc gaaggctacg tccaggagcg caccatcttc 300
ttcaaggacg acggcaacta caagaccgc gccgaggtga agttcgaggg cgacaccctg 360
gtgaaccgca tcgagctgaa gggcatcgac ttcaaggagg acggcaacat cctggggcac 420
aagctggagt acaactacaa cagccacaac gtctatatca tggccgacaa gcagaagaac 480
ggcatcaagg tgaacttcaa gatccgccac aacatcgagg acggcagcgt gcagctcgcc 540
gaccactacc agcagaacac ccccatcggc gacggccccg tgctgctgcc cgacaaccac 600
tacctgagca cccagtcgc cctgagcaaa gaccccaacg agaagcgcga tcacatggtc 660
ctgctggagt tcgtgaccgc cgccgggac actctcggca tggacgagct gtacaagtaa 720

```

<210> 11

<211> 1809

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial

Sequence: beta-glucuronidase encoding region

<400> 11

```

atggtccgtc ctgtagaaac cccaacccgt gaaatcaaaa aactcgacgg cctgtgggca 60
ttcagtcctgg atcgcgaaaa ctgtggaatt gatcagcgtt ggtgggaaag cgcgttacaa 120
gaaagccggg caattgctgt gccaggcagt ttaacgatac agttcgccga tgcagatatt 180
cgtaattatg cgggcaacgt ctggtatcag cggaagtct ttataccgaa aggttgggca 240
ggccagcgtg tcgtgctgctg tttcgatgct gtcactcatt acggcaaagt gtgggtcaat 300
aatcaggaag tgatggagca tcagggcggc tatacgccat ttgaagccga tgtcacgccg 360
tatgttattg ccgggaaaaag tgtacgtatc accgtttgtg tgaacaacga actgaactgg 420
cagactatcc cgccgggaat ggtgattacc gacgaaaacg gcaagaaaaa gcagtcttac 480
ttccatgatt tctttaacta tgccggaatc catcgacgct taatgctcta caccacgccg 540
aacacctggg tggacgatat caccgtggtg acgcatgtcg cgcaagactg taaccacgccg 600
tctgttgact ggcaggtggt ggccaatggt gatgtcagcg ttgaactgct tgatgcggat 660
caacaggtgg ttgcaactgg acaaggcact agcgggactt tgcaagtggg gaatccgcac 720
ctctggcaac cgggtgaagg ttatctctat gaactgtgctg tcacagccaa aagccagaca 780
gagtgatgata tctaccgctc tcgctcggc atccggtcag tggcagtga gggcgaaacg 840
ttcctgatta accacaaacc gttctacttt actggctttg gtcgtcatga agatgcggac 900
ttacgtggca aaggattcga taacgtgctg atggtgcacg accacgcatt aatggactgg 960
attggggcca actcctaccg tacctcgcat tacccttacg ctgaagagat gctcgactgg 1020
gcagatgaac atggcatcgt ggtgattgat gaaactgctg ctgtcggtt taacctctct 1080
ttaggcattg gtttcgaagc gggcaacaag ccgaaagaac tgtacagcga agaggcagtc 1140
aacggggaaa ctcagcaagc gcacttacag gcgattaaag agctgatagc gcgtgacaaa 1200
aaccacccaa gcgtggtgat gtggagtatt gccaacgaac cggataccccg tccgcaagtg 1260
cacgggaata tttcgccact ggcggaagca acgcgtaaac tcgacccgac gcgtccgatac 1320
acctgcgtca atgtaatggt ctgcgacgct cacaccgata ccatcagcga tctctttgat 1380
gtgctgtgcc tgaaccgtta ttacggatgg tatgtccaaa gcggcgattt ggaaacggca 1440
gagaaggtagc tggaaaaaga acttctggcc tggcaggaga aactgcatca gccgattatc 1500
atcaccgaat acggcgtgga tacgttagcc gggctgcact caatgtacac cgacatgtgg 1560
agtgaagagt atcagtgtgc atggctggat atgtatcacc gcgtctttga tcgctgcagc 1620
gccgtcgtcg gtgaacaggt atggaatttc gccgattttg cgacctcgca aggcattattg 1680
cgcgttggcg gtaacaagaa agggatcttc actcgcgacc gcaaaccgaa gtcggcggtc 1740
ttctgctgc aaaaacgctg gactggcatg aacttcggtg aaaaaccgca gcaggaggc 1800
aaacaatga 1809

```

<210> 12

<211> 411

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
part of the region of a TMV-U2 variant comprising

the origin of assembly

<400> 12

```
ccctcgccaa ttgaactcac tgaaaaagtt gttgatgagt tcgtagatga agtaccgatg 60
gctgtgaaac tcgaaagggt cgggaaaaca aaaaagagag tggtaggtaa taatgttaat 120
aataagaaaa taaataatag tggtaagaag ggtttgaaag ttgaggaaat tgaggataat 180
gtaagtgatg acgagtcctat cgcgtcatcg agtacgtttt aatcaatatg ccttatacaa 240
tcaactctcc gagccaattt gtttacttaa gttccgctta tgcagatcct gtgcagctga 300
tcaatctgtg taaaaatgca ttaggttaacc agtttcaaac gcaacaagct aggacaacag 360
tccaacagca atttgcggtat gcctggaaac ctgtgcctag tatgacagtg a 411
```

<210> 13

<211> 198

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
STMV leader region

<400> 13

```
agtaaaactt accaatcaaa agacctaac aacaggactg tcgtgggtcat ttatgctggt 60
gggggacata gggggaaaac atattgcctt ctcttacaag aggccttcag tcgccataat 120
tacttggcgc ccaattttgg gtttcagttg ctgtttccag ctatggggag aggtaagggt 180
aaaccaaacc gtaaatcg 198
```

<210> 14

<211> 455

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
STMV trailer region

<400> 14

```
gacaagtcgc cttgggttatt tcgtgttggt ttaactgaac ctgcacataa gcctttttgga 60
tcgaagggtta aacgatccgc tcctcgcttg agcttgaggc ggcgtatctc ttatgtcaac 120
agagacactt tgggtctatg ttgtataaca atagatagac tcccgtttgc aagattaggg 180
ttaacagatc ttgccgttag tctgggttagc gcgtaaccgg ccttgattta tggaatagat 240
ccattgtcca atggcctttgc caatggaacg ccgacgtggc tgtataatac gtcgttgaca 300
agtacgaaat ctgtttagtg tttttccctc cacttaaadc gaagggtttt gttttggtct 360
tcccgaacgc atacgttagt gtgactaccg ttgttcgaaa caagtaaaac aggaaggggg 420
ttcgaatccc tccctaaccg cgggtaagcg gccc 455
```

<210> 15

<211> 1971

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: cDNA copy of
part of the genome of a TMV-U1 variant, comprising
MP and CP genes

<400> 15

```

ggaaacactg tgattatagc tgcattgttg gctcgtatgc ttccgatgga gaaaataatc 60
aaaggagcct tttgtggtga cgatagtctg ctgtacttcc caaaggggtg tgagtttccg 120
gatgtgcaac actccgcgaa tcttatgtgg aattttgaag caaaactgtt taaaaaacag 180
tatggatact tttgcggaag gtatgtaata catcacgaca gaggatgcat tgtgtattac 240
gatccccata agttgatctc gaaacttggg gctaaacaca tcaaggattg ggaacacttg 300
gaggagttca gaaggtctct ttgtgatgtt gctgtttcgt tgaacaattg tgcgtattac 360
acacagttgg acgacgctgt atgggaggtt cataagaccg cccctccagg ttcgtttgtt 420
tataaaagtc tgggtgaagta tttgtctgat aaagttcttt ttagaagttt gtttatagat 480
ggctctagtt gttaaaggaa aagtgaatat caatgagttt atcgacctga caaaaatgga 540
gaagatctta ccgtcgatgt ttaccctgtt aaagagtgtc atgtgttcca aagttgataa 600
aataatggtt catgagaatg agtcattgtc agaggtaaac cttctcaaag gagttaagct 660
tattgatagt ggatacgtct gtttagccgg ttgtgtcgtc acgggcgagt ggaacttgcc 720
tgacaattgc agaggaggtg tgagcgtgtg tctggtggac aaaaggatgg aaagagccga 780
cgaggccact ctcggtatctt actacacagc agctgcaaag aaaagatttc agttcaaggt 840
cgttcccaat tatgtataaa ccaccagga cgcgatgaaa aacgtctggc aagttttagt 900
caatattaga aatgtaaaga tgtcagcggg tttctgtcgg ctttctctgg agtttgtgtc 960
ggtgtgtatc gtttatagaa ataataaaa attaggtttg agagagaaga tcacaagtgt 1020
gagagatgga gggcccatgg aacttacaga agaagttgtt gatgagttca tggaagatgt 1080
ccctatgtca atcaggcttg caaagtttcg atctcgaacc ggaaaaaaga gtgatgtccg 1140
taaagggaaa attagtagta gtgatcggtc agcgccgaac aagaactata gaaatgttaa 1200
ggattttgga ggaatgagtt ttaaaaagaa taatttaatc gatgatgatt cggagactac 1260
tgtcgccgaa tcggattcgt tttaaatatg tcttacagta tcaactacc atctcagttc 1320
gtgttcttgt cagcagcgtg ggccgaccca atagagttaa ttaatttatg tactaatgcc 1380
ttaggaaatc agtttcaaac acaacaagct cgaactgtcg ttcaaagaca attcagtgag 1440
gtgtggaaac cttcaccaca agtgactgtt aggttccctg acagtgactt taagggtgtac 1500
aggtacaatg cggattatga cccgctagtc acagcactgt taggtgcatt tgacactaga 1560
aatagaataa tagaagttga aaatcaggcg aacccccaaa ctgccgaaac gttagatgct 1620
actcgtagag tagacgacgc aacggtggcc ataaggagcg ctataaataa tttagtagta 1680
gaattgatca gaggaaccgg atcttataat cggagctctt tcgagagctc ttctgggttg 1740
gtttggaact ctggtcctgc aacttgaggt agtcaagatg cataataaat aacggattgt 1800
gtccgtaatc acacgtggtg cgtacgataa cgcatagtgt tttccctcc acttaaatcg 1860
aagggttggt tcttggtatc cgcggtgcaa atgtatatgg tcatataca tccgcaggca 1920
cgtaataaag cgaggggttc gaatcccccc gttacccccg gtaggggccc a 1971

```